



Use of Health Savings Accounts Among US Adults Enrolled in High-Deductible Health Plans

Jeffrey T. Kullgren, MD, MS, MPH^{1,2,3,4,5}; Elizabeth Q. Cliff, PhD⁶; Christopher Krenz, BA⁵ ;[et al](#)

[» Author Affiliations](#)
[| Article Information](#)



JAMA Netw Open

Published Online: July 17, 2020

2020;3;(7):e2011014.

doi:10.1001/jamanetworkopen.2020.11014



Key Points

Question How are US adults who are enrolled in a high-deductible health plan using health savings accounts to save for health care?

Findings In this national survey of 1637 respondents, approximately 1 in 3 adults enrolled in a high-deductible health plan did not have a health savings account. Among those with a health savings account, most had not contributed money into it in the last year, and less education and health insurance literacy were associated with not having made contributions.

Meaning These findings suggest that few US adults enrolled in high-deductible health plans are using health savings accounts to save for health care, and targeted interventions could enhance uptake of and contributions to health savings accounts.

Abstract

Importance Health savings accounts (HSAs) can be used by enrollees in high-deductible health plans (HDHPs) to save for health care expenses before taxes. Expansion of and encouraging contributions to HSAs have been centerpieces of recent federal legislation. Little is known about how US residents who may be eligible for HSAs are using them to save for health care.

Objective To determine which patients who may be eligible for an HSA do not have one and what decisions patients with HSAs make about contributing to them.

months were eligible to participate. Data were collected from August 26 to September 19, 2016, and analyzed from November 1, 2019, to April 30, 2020.

Main Outcomes and Measures Prevalence of not having an HSA or not making HSA contributions in the last 12 months and reasons for not making the HSA contributions.

Results Based on data from 1637 individuals (American Association of Public Opinion Research response rate 4, 54.8%), half (50.6% [95% CI, 47.7%-53.6%]) of US adults in HDHPs were female, and most were aged 36 to 51 (35.7% [95% CI, 32.8%-38.6%]) or 52 to 64 (36.8% [95% CI, 34.1%-39.5%]) years. Approximately 1 in 3 (32.5% [95% CI, 29.8%-35.3%]) did not have an HSA. Those who obtained their health insurance through an exchange were more likely to lack an HSA (70.3% [95% CI, 61.9%-78.6%]) than those who worked for an employer that offered only 1 health insurance plan (36.5% [95% CI, 30.9%-42.1%]; $P < .001$). More than half of individuals with an HSA (55.0% [95% CI, 51.1%-58.8%]) had not contributed money into it in the last 12 months. Among HDHP enrollees with an HSA, those with at least a master's degree (46.1% [95% CI, 38.3%-53.9%]; $P = .02$) or a high level of health insurance literacy (47.3% [95% CI, 40.7%-54.0%]; $P = .03$) were less likely to have made no HAS contributions. Common reasons for not contributing to an HSA included not considering it (36.8% [95% CI, 30.8%-42.8%]) and being unable to afford saving for health care (31.9% [95% CI, 26.2%-37.6%]).

Conclusions and Relevance These findings suggest that many US adults enrolled in an HDHP lack an HSA, and few with an HSA saved for health care in the last year. Targeted interventions should be explored by employers, health plans, and health systems to encourage HSA uptake and contributions among individuals who could benefit from their use.

Introduction

Health savings accounts (HSAs) can be used by US residents who are enrolled in a high-deductible health plan (HDHP) to accrue tax-free savings for health care expenses, with the policy goals of lessening patients' cost burdens and encouraging health care choices that are based on value. The proportion of privately insured adults who are enrolled in an HDHP increased from 25.3% in 2010 to 40.0% in 2016.¹ The tax-advantaged saving opportunities of HSAs could help facilitate use of needed health care services² among this growing number of HDHP enrollees, especially those who pay federal income tax, as well as preempt the cost-related access barriers and financial burdens that many experience,³⁻⁶ particularly those with lower income levels^{6,7} or a chronic condition.⁸⁻¹⁰

Beyond the potential benefits for patients who are enrolled in an HDHP,^{1,11} HSAs have been a frequent focus of recent federal health reform proposals. For example, in the 116th US Congress, at least 23 separate bills have been introduced to expand eligibility for HSAs to more than just HDHP enrollees, increase HSA contribution limits, or broaden the list of services that can be paid for with funds from an HSA.¹² Expanding use of HSAs was also part of the president's federal budgets for fiscal years 2020¹³ and 2021.¹⁴

Previous analyses have suggested that many individuals with HSAs could be saving more for health care,¹⁵ and that larger HSA balances are associated with more use of health care services.² Much less is known about which individuals who may be eligible to have an HSA lack one¹⁶ or how individuals with HSAs make choices about their health care savings.¹⁵ This information is critical to the success of policy reforms to expand uptake of and

develop targeted interventions to encourage use of HSAs as 1 strategy to help more US residents navigate the rising costs of health care.

Methods

Data

This study was deemed exempt from review by the University of Michigan Medical School institutional review board owing to the use of deidentified survey data. This survey study followed the American Association for Public Opinion Research (AAPOR) reporting guideline.

From August 26 to September 19, 2016, we surveyed 1637 participants in the GfK KnowledgePanel who had been enrolled in an HDHP (defined as a private insurance plan with a deductible of at least \$1300 for an individual or \$2600 for a family) for at least 12 months, spoke English, and were aged 18 to 64 years. The GfK KnowledgePanel is an online survey panel with approximately 55 000 US adults constructed and weighted to be representative of the US population. The GfK achieves this representativeness through address-based sampling and provision of computers and internet access to participants who lack them. People with common chronic conditions (defined as anxiety, depression, asthma, chronic obstructive pulmonary disease, heart disease, type 1 or 2 diabetes, or hypertension) were oversampled because they often face greater financial burdens or forgo needed health care services when facing high cost sharing.^{3,6,8-10} Participants provided oral or written consent to GfK to participate in the KnowledgePanel and to receive periodic survey invitations from GfK. Additional details about the GfK KnowledgePanel are available in eMethods in the [Supplement](#). The American Association for Public Opinion Research survey response rate 4, defined as the number of completed plus partially completed surveys out of the number of individuals who were known to be eligible plus an estimate of the proportion of individuals with unknown eligibility who were eligible,¹⁷ was 54.8% (eFigure in the [Supplement](#)).

Survey Measures

Respondents were asked if they had an HSA, defined in the National Health Interview Survey as "a special account or fund that can be used to pay for medical expenses" that are "sometimes referred to as health savings accounts (HSAs), health reimbursement accounts (HRAs), personal care accounts, personal medical funds, or choice funds, and are different from flexible spending accounts."¹ (p10) Respondents were also asked if they saved for health care expenses in the last 12 months and, if so, which savings vehicles they used and how much they saved. Respondents who had not saved for health care in the last 12 months (in an HSA or any other savings vehicle) were asked to choose reasons why they had not saved from a list and could also write in reasons. Other survey questions asked about health status and source of insurance coverage, as well as factors that could potentially mediate health care savings, such as levels of consumer engagement,¹⁸ financial literacy,¹⁹ and health insurance literacy.²⁰ To optimize participant understanding and internal validity, all new survey measures were refined through iterative cognitive interviews with 17 US adults in HDHPs with varying age, sex, sources of health insurance, geography, health literacy, and health care experience. New measures were then finalized after additional testing among 100 Amazon Mechanical Turk participants²¹ and 60 GfK KnowledgePanel participants. Survey responses were supplemented with demographic data collected by GfK, including participants' self-report of their race and ethnicity. The full survey instrument is available in the eAppendix in the [Supplement](#).

Statistical Analysis



Sections



PDF



Share

the March 2016 Current Population Survey and the 2015 National Health Interview Survey. To assess potential nonresponse bias, characteristics of survey respondents and nonrespondents were compared (eTable 1 in the [Supplement](#)), and the weighted sample was compared with weighted characteristics of nonelderly National Health Interview Survey participants enrolled in an HDHP (eTable 2 in the [Supplement](#)).

Weighted multivariable logistic regression was used to estimate the association between not having an HSA and demographic characteristics, health status, source of insurance, having a chronic condition, level of consumer engagement, level of health insurance literacy, and level of financial literacy. Among the subpopulation of individuals with an HSA, weighted multivariable logistic regression was used to estimate the association of not having made any HSA contributions in the last 12 months and these same independent variables. All independent variables were operationalized as mutually exclusive categories. Scores from scales used to measure potential mediators of saving (ie, consumer engagement, financial literacy, and health insurance literacy) were grouped into tertiles.²²⁻²⁴ Using the estimated coefficients from each regression model, we report marginal estimates of the adjusted prevalence of each outcome as a function of each independent variable.²⁵

Analyses were conducted from November 1, 2019, to April 30, 2020. Stata, version 15.1 (StataCorp LLC), was used for all analyses. Variance estimates were computed using Taylor series linearization to reflect the variability of the survey weights. Two-tailed *P* values were obtained from logistic regression models, and *P* < .05 indicated significance.

Results

Population Characteristics

Based on weighted data from 1637 survey respondents, we estimate that half (50.6% [95% CI, 47.7%-53.6%]) of US adults in HDHPs were female, and most were aged 36 to 51 (35.7% [95% CI, 32.8%-38.6%]) or 52 to 64 (36.8% [95% CI, 34.1%-39.5%]) years. Most were employed (83.8% [95% CI, 81.6%-86.0%]) and received their health insurance through an employer (84.6% [95% CI, 82.6%-86.5%]). Nearly half (42.4% [95% CI, 39.6%-45.3%]) had a chronic condition. Additional characteristics of the study population have been previously published²²⁻²⁴ and are available in eTable 3 in the [Supplement](#). Compared with survey nonrespondents, survey respondents were older (aged 52-64 years, 48.6% [95% CI, 46.2%-51.0%] vs 32.9% [95% CI, 31.6%-34.2%]), more likely to be men (53.9% [95% CI, 51.5%-56.3%] vs 43.5% [42.1%-44.9%]) and white (81.6% [95% CI, 79.7%-83.5%] vs 69.0% [95% CI, 67.7%-70.3%]), and had higher levels of education (bachelor's degree, 34.1% [95% CI, 31.9%-36.4%] vs 28.8% [95% CI, 27.6%-30.1%]) and income (≥400% of the federal poverty line, 53.7% [95% CI, 51.3%-56.1%] vs 46.1% [95% CI, 44.7%-47.5%]) (eTable 1 in the [Supplement](#)).

Prevalence of Lacking an HSA Overall and Among Subgroups


Among US adults with an HDHP, an estimated 32.5% (95% CI, 29.8%-35.3%) did not have an HSA, 58.4% (95% CI, 55.4%-61.3%) had an HSA, and 9.1% (95% CI, 7.5%-11.0%) did not know whether they had an HSA or did not complete the survey question about having an HSA. Compared with HDHP enrollees who worked for an employer that offered only 1 plan (36.5% [95% CI, 30.9%-42.1%]), those who worked for an employer that offered multiple plans were less likely to lack an HSA (21.5% [95% CI, 18.3%-24.7%]; *P* < .001), and those who obtained coverage through an exchange were more likely to lack an HSA (70.3% [95% CI, 61.9%-78.6%]; *P* < .001) ([Table 1](#)).

Individuals with a master's degree (22.1% [95% CI, 17.0%-27.3%]) were significantly less likely than those with a

compared with those with higher incomes or for those with vs without chronic conditions. All estimated model parameters are shown in eTable 4 in the [Supplement](#).



Table 1. Marginal Estimates of the Adjusted Prevalence of Reporting No HSA Among US Adults in HDHPs

 Marginal Estimates of the Adjusted Prevalence of Reporting No HSA Among US Adults in HDHPs


[Go to Figure in Article](#)

Prevalence of Making No HSA Contributions Overall and Among Subgroups

Among HDHP enrollees with an HSA, more than half (55.0% [95% CI, 51.1%-58.8%]) had not contributed any money into it in the last 12 months ([Figure 1](#)). Compared with HDHP enrollees with an HSA who had no more than a high school education (62.6% [95% CI, 52.4%-72.8%]), those who had a bachelor's degree (49.5% [95% CI, 43.2%-55.7%]; $P = .04$) or at least a master's degree (46.1% [95% CI, 38.3%-53.9%]; $P = .02$) were less likely to have made no HSA contributions in the last 12 months ([Table 2](#)). Individuals with an HSA who had a high level of health insurance literacy (47.3% [95% CI, 40.7%-54.0%]) were significantly less likely to have made no HSA contributions in the last 12 months than those with a low level of health insurance literacy (58.6% [95% CI, 51.5%-65.6%]; $P = .03$). Compared with HDHP enrollees with an HSA who had obtained their coverage from an employer that offered only 1 insurance plan (61.0% [95% CI, 53.2%-68.7%]), those who had obtained their coverage from an exchange (30.9% [95% CI, 6.9%-54.9%]) were less likely to have made no HSA contributions in the last 12 months ($P = .03$). There were no statistically significant differences in the adjusted prevalence of having made no HSA contributions in the last 12 months by income level or the presence of a chronic condition. All estimated model parameters are shown in eTable 5 in the [Supplement](#).



Figure 1. Health Care Savings in the Last Year Among US Adults in High-Deductible Health Plans (HDHPs) Who Had a Health Savings Account (HSA)

 Health Care Savings in the Last Year Among US Adults in High-Deductible Health Plans (HDHPs) Who Had a Health Savings Account (HSA)

[Go to Figure in Article](#)

Includes 929 respondents with proportions weighted for US population. Error bars indicate 95% CIs.



Table 2. Marginal Estimates of the Adjusted Prevalence of Making No HSA Contributions in the Last Year Among US Adults in HDHPs Who Had an HSA

Reasons for Not Contributing to HSAs

The most common reasons for not saving in the last 12 months among people with an HSA were that they did not perceive a need for health care services (44.9% [95% CI, 38.7%-51.1%]) or that they already had sufficient savings to cover health care expenses (40.2% [95% CI, 34.2%-46.3%]) (**Figure 2**). Other common reasons for not contributing to an HSA included not considering it (36.8% [95% CI, 30.8%-42.8%]) and being unable to afford saving for health care (31.9% [95% CI, 26.2%-37.6%]).



Figure 2. Reasons for Not Saving for Health Care Among US Adults in High-Deductible Health Plans (HDHPs) Who Had a Health Savings Account (HSA)

 [Reasons for Not Saving for Health Care Among US Adults in High-Deductible Health Plans \(HDHPs\) Who Had a Health Savings Account \(HSA\)](#)

[Go to Figure in Article](#)


Includes 366 respondents with proportions weighted for US population. Error bars indicate 95% CIs.

Savings Among HDHP Enrollees Who Had an HSA

Of those with an HSA who did contribute money into their account in the last 12 months, the most frequent level of contribution was \$2000 or more (46.7% [95% CI, 41.2%-52.2%]). For most individuals who contributed to their HSA (59.4% [95% CI, 54.5%-64.2%]), the HSA was the only place they saved for health care (**Figure 3**). Another 21.4% (95% CI, 17.6%-25.8%) of those who put money into their HSA used another vehicle to save for health care (eg, a flexible spending account or bank account) in addition to their HSA.



Figure 3. Savings Vehicles Used by US Adults in High-Deductible Health Plans (HDHPs) Who Had a Health Savings Account (HSA) and Saved for Heath Care

 [Savings Vehicles Used by US Adults in High-Deductible Health Plans \(HDHPs\) Who Had a Health Savings Account \(HSA\) and Saved for Heath Care](#)

[Go to Figure in Article](#)

Includes 548 respondents with proportions weighted for US population. Error bars indicate 95% CIs; FSA, flexible spending account.

In this survey study, few US adults who were enrolled in HDHPs were using HSAs to save for health care expenses, despite the cost-related barriers to access and financial burdens experienced by the growing number of HDHP enrollees.^{3-7,10} Previous analyses have examined levels of HSA contributions and balances¹⁵ and their relation to the use of health care services² among people who have HSAs. This study is among the first, to our knowledge, to examine how US adults who may be eligible for HSAs are using these accounts to save for future health care expenses and identifies opportunities for new interventions that employers, health plans, and health systems could implement to encourage use of HSAs among individuals who are eligible to have them and could benefit from their use.

Approximately 1 in 3 US adults who were enrolled in an HDHP reported not having an HSA. Some HDHP enrollees were more likely than others to lack an HSA, particularly individuals who obtained their insurance through an exchange. The relatively high rate of exchange plans lacking an HSA may be because, although the mean deductible for a federal health insurance Marketplace individual plan was \$5316 during the 2020 open enrollment period, just 7% of Marketplace plans chosen for 2020 were qualified for linkage to an HSA.²⁶ This low rate of HSA eligibility could be owing to some Marketplace HDHPs covering services before the deductible in ways that are not currently allowed in HSA-eligible plans.²⁷ Consequently, a policy option for increasing uptake of HSAs in Marketplace plans would be to allow more flexibility in what types of HDHPs qualify for linkage to an HSA.

Nevertheless, our results also suggest that legislative efforts to expand eligibility for HSAs may not have their full intended effects without concomitant efforts to support eligible individuals to take advantage of their eligibility and facilitate their acquisition of an HSA. In the case of HDHPs obtained through an exchange, exchanges could better highlight the benefits of HSAs to encourage uptake of HSA-eligible plans at the time of enrollment. Alternatively, policy reforms could require that HDHPs offered on an exchange be eligible for and linked to an HSA. After enrollment in an HDHP, employers, health plans, and health systems could target messaging to HDHP enrollees to encourage acquisition of an HSA as a strategy to help manage the high cost-sharing of these plans. Although we collected data on reasons for not contributing to an HSA among those who had one, we were not able to collect data on why individuals without an HSA had not acquired one. This is an important area for future research, which would provide valuable information on whether these interventions, or other strategies, can help facilitate uptake of HSAs among individuals who are eligible to have them.

Most individuals who reported having an HSA had not contributed any savings into their account in the previous 12 months. Although some of these individuals may have had a positive balance in their account from previous contributions,¹⁵ already having sufficient savings was only cited by 40% of individuals as a reason for not having saved for health care in the last 12 months. This result suggests that current legislative efforts to increase the HSA contribution limit would likely have little effect on health care savings for the vast majority of people with HSAs and that other factors may be driving consumers' decisions about HSA savings. Indeed, additional common reasons for not having saved for health care included not having considered saving and being unable to afford saving. To prompt consideration of contributing to an HSA, health plans and health systems could periodically target educational messaging to HDHP enrollees who have an HSA. Among individuals who have an employer-sponsored HDHP with an HSA but have not contributed to it, employers could facilitate more HSA savings through interventions that have been successful at increasing retirement savings in workplaces, such as default contributions to HSAs, employer matching of employee HSA contributions, or committing future wage increases to savings.²⁸



Approximately 1 in 5 individuals with an HSA who saved for health care in the last 12 months reported saving in a non-HSA vehicle along with their HSA. For individuals who have not reached their annual HSA contribution limit, this may be financially inferior to the triple tax advantage of saving in an HSA (ie, HSA contributions are not taxed, HSA earnings are not taxed, and qualified HSA withdrawals are not taxed). Another 20% of people with an HSA reported saving for health care only in a non-HSA vehicle and could similarly be missing opportunities to reduce their tax burdens. Together, these patterns suggest that employers and health plans have potentially important roles to play in helping individuals with HSAs better understand these accounts and make financially advantageous decisions about their saving for health care.

Although people with lower incomes or chronic conditions can be especially vulnerable to cost-related access barriers when enrolled in an HDHP,⁶⁻¹⁰ individuals in these subgroups were not less likely than others to lack an HSA. Similarly, among individuals who had an HSA, those with lower incomes or chronic conditions were not any less likely than others to have made no contributions to their HSA in the last year. Efforts by employers, health plans, and health systems to encourage uptake and use of HSAs should particularly focus on these subgroups of HDHP enrollees who could potentially benefit from the use of HSAs as a strategy to lessen their known cost-related access barriers and financial burdens.

Limitations

These data are self-reported and thus subject to recall bias. To limit such bias, existing survey measures were used where possible, and new measures were developed through extensive cognitive interviews.²² To classify people as having an HSA, an existing National Health Interview Survey measure was used that relied on individuals' knowledge, because it was not feasible to collect details about their accounts. Although this approach is consistent with many other studies,^{1,29,30} it means that we were unable to confirm that participants' HSAs were Internal Revenue Service-qualified HSAs. If someone reported having multiple savings vehicles, including an HSA, we assumed that all reported savings for health care went into their HSA, which would bias reported HSA savings upward. To check the sensitivity of our results to this assumption, we analyzed levels of savings among those who only saved in an HSA and found that this did not change our main conclusions. We did not inquire about reasons for not enrolling in an HSA or, among people with an HSA, balances or expenditures for their HSA, and thus were unable to examine these issues.

Conclusions

The results of this survey study suggest that few US adults enrolled in HDHPs are using HSAs to save for their health care expenses. As enrollment in HDHPs and policy interest in expanding use of HSAs both continue to grow, employers, health plans, and health systems should explore targeted interventions to encourage HSA uptake and contributions among individuals whose use of an HSA could potentially improve the affordability of needed health care services.

Article Information

Accepted for Publication: May 7, 2020.

Published: July 17, 2020. doi:10.1001/jamanetworkopen.2020.11014

Corresponding Author: Jeffrey T. Kullgren, MD, MS, MPH, Veterans Affairs Center for Clinical Management Research, Veterans Affairs Ann Arbor Healthcare System, PO Box 130170, Ann Arbor, MI 48113 (jkullgre@med.umich.edu).

Author Contributions: Dr Kullgren had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Kullgren, West, Levy, Fendrick, Fagerlin.

Acquisition, analysis, or interpretation of data: Kullgren, Cliff, Krenz, West, Levy.

Drafting of the manuscript: Kullgren, Cliff, West.

Critical revision of the manuscript for important intellectual content: Kullgren, Krenz, West, Levy, Fendrick, Fagerlin.

Statistical analysis: Kullgren, Cliff, West, Levy.

Obtained funding: Kullgren, West.

Administrative, technical, or material support: Cliff, Krenz, West.

Supervision: Kullgren, Levy.

Conflict of Interest Disclosures: Dr Kullgren reported receiving grants from the Robert Wood Johnson Foundation during the conduct of the study and grants from HealthWell Foundation and personal fees from SeeChange Health, HealthMine, Kaiser Permanente Washington Health Research Institute, Robert Wood Johnson Foundation, AbilTo, Inc, Kansas City Area Life Sciences Institute, and the American Diabetes Association outside the submitted work. Dr Cliff reported receiving grants from Robert Wood Johnson Foundation during the conduct of the study and an ABIM Foundation grant outside the conduct of the study. Mr Krenz reported receiving grants from Robert Wood Johnson Foundation during the conduct of the study. Dr Fendrick reported consulting for AbbVie, Inc, Amgen, Inc, Centivo, Community Oncology Association, Covered California, EmblemHealth, Exact Sciences Corp, Freedman Health System, GRAIL, Harvard University, Health & Wellness Innovations, Health[at]Scale Technologies, HealthCorum, MedZed LLC, Merck & Co, Montana Health Cooperative, Penguin Pay, Phathom Pharmaceuticals, Risalto, Sempre Health, the State of Minnesota, US Department of Defense, Virginia Center for Health Innovation, Wellth, Yale-New Haven Health System, and Zansors; having an equity interest in Health & Wellness Innovations, Health[at]Scale Technologies, Sempre Health, Wellth, and Zansors; receiving research support from the Agency for Healthcare Research and Quality, Boehringer-Ingelheim, the Gary and Mary West Health Policy Center, Laura and John Arnold Foundation, National Pharmaceutical Council, Patient-Centered Outcomes Research Institute, Pharmaceutical Research and Manufacturers of America, and the Michigan Department of Health and Human Services; and serving as coeditor of the *American Journal of Managed Care*, a member of the Medicare Evidence Development and Coverage Advisory Committee, and a partner in V-BID Health, LLC. No other disclosures were reported.

Funding/Support: This study was supported by grant 73054 from the Robert Wood Johnson Foundation; by the Department of Veterans Affairs, Veterans Health Administration, Health Services Research and Development

Role of the Funder/Sponsor: The sponsors had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Disclaimer: The views expressed in this article are those of the authors and do not necessarily reflect the view of the Robert Wood Johnson Foundation or the position or policy of the Department of Veterans Affairs or the US government.

Meeting Presentation: This paper was presented at the 2019 Society of General Internal Medicine Annual Meeting (May 9, 2019; Washington, DC) and the 2019 AcademyHealth Annual Research Meeting (June 2, 2019; Washington, DC).

References

1. Cohen RA, Martinez ME, Zammitti EP; National Center for Health Statistics. Health insurance coverage: early release of estimates from the National Health Interview Survey, January-March 2016. Released September 2016. Accessed November 1, 2019.
<https://www.cdc.gov/nchs/data/nhis/earlyrelease/insur201609.pdf>
2. Fronstin P, Roebuck MC. Do accumulating HSA balances affect use of health care services and spending? Employee Benefit Research Institute. Posted May 23, 2019. Accessed February 6, 2020.
<https://www.ebri.org/health/publications/issue-briefs/content/do-accumulating-hsa-balances-affect-use-of-health-care-services-and-spending>
3. Dixon A, Greene J, Hibbard J. Do consumer-directed health plans drive change in enrollees' health care behavior? *Health Aff (Millwood)*. 2008;27(4):1120-1131. doi:[10.1377/hlthaff.27.4.1120](https://doi.org/10.1377/hlthaff.27.4.1120)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
4. Hibbard JH, Greene J, Tusler M. Does enrollment in a CDHP stimulate cost-effective utilization? *Med Care Res Rev*. 2008;65(4):437-449. doi:[10.1177/1077558708316686](https://doi.org/10.1177/1077558708316686)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
5. Greene J, Hibbard J, Murray JF, Teutsch SM, Berger ML. The impact of consumer-directed health plans on prescription drug use. *Health Aff (Millwood)*. 2008;27(4):1111-1119. doi:[10.1377/hlthaff.27.4.1111](https://doi.org/10.1377/hlthaff.27.4.1111)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
6. Abdus S, Keenan PS. Financial burden of employer-sponsored high-deductible health plans for low-income adults with chronic health conditions. *JAMA Intern Med*. 2018;178(12):1706-1708. doi:[10.1001/jamainternmed.2018.4706](https://doi.org/10.1001/jamainternmed.2018.4706)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
7. Kullgren JT, Galbraith AA, Hinrichsen VL, et al. Health care use and decision making among lower-income families in high-deductible health plans. *Arch Intern Med*. 2010;170(21):1918-1925. doi:[10.1001/archinternmed.2010.428](https://doi.org/10.1001/archinternmed.2010.428)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
8. Galbraith AA, Ross-Degnan D, Soumerai SB, Rosenthal MB, Gay C, Lieu TA. Nearly half of families in

9. Galbraith AA, Soumerai SB, Ross-Degnan D, Rosenthal MB, Gay C, Lieu TA. Delayed and forgone care for families with chronic conditions in high-deductible health plans. *J Gen Intern Med*. 2012;27(9):1105-1111. doi:[10.1007/s11606-011-1970-8](#)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
10. Segel JE, Kullgren JT. Health insurance deductibles and their associations with out-of-pocket spending and affordability barriers among US adults with chronic conditions. *JAMA Intern Med*. 2017;177(3):433-436. doi:[10.1001/jamainternmed.2016.8419](#)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
11. Fronstin P. Enrollment in HSA-eligible health plans: slow and steady growth continued into 2018. Employee Benefit Research Institute. Posted March 28, 2019. Accessed February 6, 2020.
<https://www.ebri.org/health/publications/issue-briefs/content/enrollment-in-hsa-eligible-health-plans-slow-and-steady-growth-continued-into-2018>
12. Congress.gov. Library of Congress. Updated June 15, 2020. Accessed December 17, 2019.
<https://www.congress.gov/>
13. Budget of the US Government. Fiscal Year 2020. Posted March 11, 2019. Accessed April 1, 2020.
<https://www.whitehouse.gov/wp-content/uploads/2019/03/budget-fy2020.pdf>
14. Budget of the US Government. Fiscal Year 2021. Posted February 10, 2020. Accessed April 1, 2020.
https://www.whitehouse.gov/wp-content/uploads/2020/02/budget_fy21.pdf
15. Fronstin P, Spiegel J. Trends in health savings account balances, contributions, distributions, and investments, 2011-2018: estimates from the EBRI HSA database. Employee Benefit Research Institute. Posted January 9, 2020. Accessed February 6, 2020. <https://www.ebri.org/health/publications/issue-briefs/content/trends-in-health-savings-account-balances-contributions-distributions-and-investments-2011-2018-estimates-from-the-ebri-hsa-database>
16. Helmchen LA, Brown DW, Lurie IZ, Lo Sasso AT. Health savings accounts: growth concentrated among high-income households and large employers. *Health Aff (Millwood)*. 2015;34(9):1594-1598. doi:[10.1377/hlthaff.2015.0480](#)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
17. American Association for Public Opinion Research. Standard definitions: final dispositions of case codes and outcome rates for surveys. 9th ed. Revised 2016. Accessed February 6, 2020.
http://www.aapor.org/AAPOR_Main/media/publications/Standard-Definitions20169theditionfinal.pdf
18. Duke CC, Lynch WD, Smith B, Winstanley J. Validity of a new patient engagement measure: the Altarum Consumer Engagement (ACE) measure. *Patient*. 2015;8(6):559-568. doi:[10.1007/s40271-015-0131-2](#)
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
19. Lusardi A, Mitchell OS. The economic importance of financial literacy: theory and evidence. National Bureau of Economic Research. Issued April 2013. Accessed July 28, 2015.
<https://www.nber.org/papers/w18952>

21. Amazon Mechanical Turk. Access a global, on-demand 24 × 7 workforce. Updated April 8, 2020. Accessed June 14, 2017. <https://worker.mturk.com/mturk/welcome>
22. Kullgren JT, Cliff BQ, Krenz CD, et al. A survey of Americans with high-deductible health plans identifies opportunities to enhance consumer behaviors. *Health Aff (Millwood)*. 2019;38(3):416-424. doi:10.1377/hlthaff.2018.05018
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
23. Kullgren JT, Cliff EQ, Krenz C, et al. Consumer behaviors among individuals enrolled in high-deductible health plans in the United States. *JAMA Intern Med*. 2018;178(3):424-426. doi:10.1001/jamainternmed.2017.6622
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
24. Cliff BQ, Krenz C, West BT, et al. Attitudes about consumer strategies among Americans in high-deductible health plans. *Med Care*. 2019;57(3):187-193. doi:10.1097/MLR.0000000000001056
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
25. Graubard BI, Korn EL. Predictive margins with survey data. *Biometrics*. 1999;55(2):652-659. doi:10.1111/j.0006-341X.1999.00652.x
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
26. Health Insurance Exchanges 2020 Open Enrollment Report. Centers for Medicare & Medicaid Services. Updated May 8, 2020. Accessed June 15, 2020. <https://www.cms.gov/files/document/4120-health-insurance-exchanges-2020-open-enrollment-report-final.pdf>
27. O'Brien E. Why 80% of Obamacare plans are ineligible for this tax break? MarketWatch. Published March 29, 2016. Accessed May 1, 2020. <https://www.marketwatch.com/story/why-80-of-obamacare-plans-are-ineligible-for-this-tax-break-2016-03-29>
28. Thaler RH, Benartzi S. Save more tomorrow: using behavioral economics to increase employee saving. *J Polit Econ*. 2004;112(S1):S164-S187. doi:10.1086/380085
[Google Scholar](#) | [Crossref](#)
29. Hageman SA, St George DMM. Health savings account ownership and financial barriers to health care: what social workers should know. *Soc Work Public Health*. 2019;34(2):176-188. doi:10.1080/19371918.2019.1575310
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
30. Zheng Z, Jemal A, Banegas MP, Han X, Yabroff KR. High-deductible health plans and cancer survivorship: what is the association with access to care and hospital emergency department use? *J Oncol Pract*. 2019;15(11):e957-e968. doi:10.1200/JOP.18.00699
[PubMed](#) | [Google Scholar](#) | [Crossref](#)

