

A Flexible Fourier Approach to Repeat Sales Price Indexes

Daniel P. McMillen, Jonathan Dombrow

First published: 17 December 2002

<https://doi.org/10.1111/1080-8620.00008>

Citations: 44

Abstract

Time periods are typically highly aggregated for repeat sales estimators because of the small number of observations available in some periods. We use a flexible Fourier expansion to account for time, which we treat as a continuous variable. Our estimator saves degrees of freedom and enables us to estimate the price index efficiently even for times with few sales. We present estimated price indexes for the City of Chicago, Cook County, and several suburbs.

Citing Literature



[Download PDF](#)

ABOUT WILEY ONLINE LIBRARY

[Privacy Policy](#)

[Terms of Use](#)

[About Cookies](#)

[Manage Cookies](#)

[Accessibility](#)

[Wiley Research DE&I Statement and Publishing Policies](#)

[Developing World Access](#)

HELP & SUPPORT

Contact Us
Training and Support
DMCA & Reporting Piracy

OPPORTUNITIES

Subscription Agents
Advertisers & Corporate Partners

CONNECT WITH WILEY

The Wiley Network
Wiley Press Room

Copyright © 1999-2024 John Wiley & Sons, Inc or related companies. All rights reserved, including rights for text and data mining and training of artificial intelligence technologies or similar technologies.

WILEY