



## Notes

To simulate symmetric stable random variables the algorithm of Chambers et al. (<u>1976</u>), encoded in GAUSS by J. Huston McCulloch, is used.

For series with less than 200 observations the maximum lag length is set to 24.

Here and elsewhere, to estimate the index of stability

the method suggested by McCulloch (1986) is used.

Related Research Data			
LAG Length Selection and the Construction of Unit Root Tests with Good Size and			
Power			
Source: Econometrica			
Simple consistent estimators of stable distribution parameters			
Source: Communications in Statistics - Simulation and Computation			
Time contractive with marked and infinite mentioned disturbances			
Source			
Unit F of the			
Trunc			
Sourc			
Time			
Sourc			
lestin			
Source			
A Met			
Source			
Critic			
Source			
Time			
Sourc			

## Related research (1)

People also read	Recommended articles	Cited by 40		
Information for	Open access	Open access		
Authors	Overview	Overview		
R&D professionals	Open journals	Open journals		
Editors	Open Select	Open Select		
Librarians	Dove Medical Pre	Dove Medical Press		
Societies	F1000Research	F1000Research		
Opportunities	Help and inform	Help and information		
Reprints and e-prints	Help and contact	Help and contact		
Advertising solutions	Newsroom	Newsroom		
Accelerated publication	All journals	All journals		
Corporate access solutions	Books	Books		

Keep up Register t by email Sigr Copyright

Registered 5 Howick Pl ×

or & Francis Group