



Q

Home ► All Journals ► Food Science & Technology ► British Poultry Science ► List of Issues ► Volume 49, Issue 1 ► Spoilage of light (PSE-like) and dark tu

British Poultry Science > Volume 49, 2008 - Issue 1

486440ViewsCrossRef citations to dateAltmetric

Original Articles

Spoilage of light (PSE-like) and dark turkey meat under aerobic or modified atmosphere package: microbial indicators and their relationship with total volatile basic nitrogen

Dr M.J. Fraqueza , M.C. Ferreira & A.S. Barreto Pages 12-20 | Accepted 31 Jul 2007, Published online: 21 Jan 2008

G Cite this article **A** https://doi.org/10.1080/00071660701821675

Sample our Bioscience Journals >> Sign in here to start your access

🖹 Full A

🔒 Repri

Abstra



evaluati

2. Brea for light colour. S 50% CO

We Care About Your Privacy

We and our 907 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting "I Accept" enables tracking technologies to support the purposes shown under "we and our partners process data to provide," whereas selecting "Reject All" or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the ["privacy preferences"] link on the bottom of the webpage [or the floating icon on the bottom-left of the webpage, if applicable]. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. <u>Here</u>

We and our partners process data to provide:

L Accept Reject All Show Purpose Jifferent rk), stablish a -N), J pH < 5.8 r dark 6 N₂ and under

aerobic conditions was evaluated for microbiological characteristics and TVB-N on d 0,

5 and 12. This evaluation was extended to include d 19 and 25 when samples were under MAP conditions.

3. The dark meat group after 12 d of storage in aerobiosis presented significantly higher plate counts of aerobic mesophilic, psychrotrophic micro-organisms and higher TVB-N than other meat colour categories. The shelf life of turkey meat under MAP was one week longer for intermediate and light colour meat (20 d) than for dark meat. TVB-N values of 20 to 30 mg NH₃/100 g turkey meat correspond to advanced spoilage stages. We proposed 14 mg NH₃/100 g as the limit of freshness acceptability for turkey meat.

4. TVB-N was an indicator of turkey meat microbial spoilage but was not a suitable early predictor for microbial spoilage and in particular for turkey meat stored under MAP conditions because counts of micro-organisms were moderately correlated (Pseudomonas spp. and Enterobacteriaceae) with this index, as they were inhibited by MAP gas mixture and storage temperature used in the present study.



Information for	Open access
Authors	Overview
R&D professionals	Open journals
Editors	Open Select
Librarians	Dove Medical Press
Societies	F1000Research
Opportunities	Help and information
Reprints and e-prints	Help and contact
Advertising solutions	Newsroom
Accelerated publication	All journals
Corporate access solutions	Books

Keep up to date

Register to receive personalised research and resources by email

🔛 Sign me u

