









References

66 Citations

Metrics

➡ Reprints & Permissions

Read this article

Share

Abstract

It is known that the substitution of cations in perovskites produce changes in the macroscopic properties of these materials. A case to study is for example the influence of 4^+ -cation B' partially substituted by other 4^+ -cation B' when ABO3 is ferroelectric and AB'O3 is paraelectric. In this work the system $BaTi_{1-x}Hf_xO3$ with $x=0.05,\,0.10,\,0.15$ and 0.20 is studied by Perturbed Angular Correlations (PAC) spectroscopy in order to get microscopic information through the electric field gradient tensor (EFG) produced by electrons close to probes. Samples were prepared by solid state reactions and characterized by x-ray powder diffraction analyses and irradiated with thermal neutrons to produce 181 Ta, the PAC probe. Two hyperfine quadrupole interactions were detected. One of them correspond to probes in sites with defects originated during the nuclear processes after neutron irradiation. The other probes are located in B sites. At RT the hyperfine parameters are analyzed in terms of Hf concentration.

Related research 1



People also read

Recommended articles

Cited by 3

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up









Accessibility





Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions Taylor and Francis Group

Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG