



Journal of Motor Behavior >

Volume 46, 2014 - [Issue 2](#)

562 | 16 | 0
Views | CrossRef citations to date | Altmetric

Research Articles

Trunk, Head, and Step Characteristics During Normal and Narrow-Based Walking Under Deteriorated Sensory Conditions

Nandini Deshpande & Fang Zhang

Pages 125-132 | Received 20 Jun 2013, Accepted 16 Dec 2013, Published online: 14 Feb 2014

🗨️ Cite this article 🔗 <https://doi.org/10.1080/00222895.2013.877416>



Sample our
Behavioral Sciences
Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

📄 Full Article

🖼️ Figures & data

📖 References

🗨️ Citations

📊 Metrics

📄 Reprints & Permissions

Read this article

🔗 Share

ABSTRACT

The ability to maintain stability in the frontal plane (medialateral direction) while walking is commonly included as a component of motor performance assessment. Postural control in the frontal plane may deteriorate faster and earlier with increasing age, compared to that in the sagittal plane (anteroposterior direction). Fifteen young (20-30 years old) and 15 older (>65 years old) healthy participants were recruited to investigate age-related differences in postural control during the normal and narrow-based walking when performed under suboptimal vestibular and lower limb somatosensory conditions achieved by galvanic stimulation and compliant surfaces, respectively. Gait speed decreased in the narrow-based walking condition, with larger decrease in the elderly (by 6%). In the elderly head roll increased with perturbed vestibular information in impaired somatosensory condition (by 40.70%). In both age groups trunk roll increased under impaired somatosensation in the narrow-based

walking condition (by 43.62%) but not in normal walking condition. Older participants adopted a more cautious strategy characterized by lower walking speed when walking on a narrow base and exhibited deteriorated integrative ability of the CNS for head control. Accurate lower limb somatosensation may play a critical role in narrow-based walking.

Keywords:

- gait
- head control
- trunk control
- step characteristics
- sensory integration

ACKNOWLEDGMENTS

The authors thank Dr. Alison Novak, Mika Yoshikawa, and Patricia Hewston for assistance with data collection.

Related research

People also read

Recommended articles

Cited by
16

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)

 Taylor and Francis Group

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG