Taylor & Francis
Online

Molecular Physics ›
An International Journal at the Interface Between Chemistry and Physics
Volume 111, 2013 - Issue 14-15: Dedicated to Martin Quack on the Occasion of his 65th Birthday

302 Views | 30 CrossRef citations to date | 0 Altmetric

Invited Article

Absolute frequency measurements of CO_2 transitions at 4.3 μm with a comb-referenced quantum cascade laser

Iacopo Galli, Saverio Bartalini, Pablo Cancio Pastor, Francesco Cappelli, Giovanni Giusfredi, Davide Mazzotti, ...show all

Pages 2041-2045 | Received 15 Jan 2013, Accepted 27 Feb 2013, Accepted author version posted online: 06 Mar 2013, Published online: 03 Apr 2013

Cite this article https://doi.org/10.1080/00268976.2013.782436

Sample our Mathematics & Statistics Journals
>> Sign in here to start your access to the latest two volumes for 14 days

Full Article Figures & data References Citations Metrics
Reprints & Permissions Read this article Share

Abstract

The infrared spectrum of the (01^11–01^10) ro-vibrational band of $^{12}C^{16}O_2$ in the range 2306–2312 cm^{-1} is investigated with saturated-absorption sub-Doppler spectroscopy. The absolute frequencies of six transitions belonging to the P-branch of this band are measured by recording their Lamb-dip features in a pump-probe detection scheme employing a room-temperature quantum cascade laser. The laser is phase-locked to a subkilohertz-linewidth difference-frequency-generated radiation source, which is referenced to an optical frequency comb synthesiser. The achieved relative uncertainties range from 1×10^{-11} to 5×10^{-11}, improving by three to four orders of magnitude the previous tabulated values for such frequencies. Moreover, thanks to this

precision level, self-pressure-shift coefficients due to collisional processes of CO_2 molecules are reported for the first time.

Keywords:

carbon dioxide | saturated-absorption spectroscopy | Lamb dip | quantum cascade laser | optical frequency comb

Acknowledgements

We wish to thank Prof. Gianfranco Di Lonardo for his valuable help in finding the most accurate calculated frequencies of our observed transitions. This work was financially supported by Ente Cassa di Risparmio di Firenze, by the Laserlab-Europe Consortium in the ALADIN project framework, by the Extreme Light Infrastructure (ELI) European project and by the Progetto Operativo Nazionale (PON) PON01_01525 ‘MONitoraggio Innovativo per le Coste e l’Ambiente marino’ (MONICA) funded by the Italian Ministry of Education, University and Research (MIUR).

Notes

aComparison with the HITRAN database [3–4]. The uncertainty reported by the original database for each transition is between 3 and 30 MHz.

bComparison with Ref. [29], corrected by the calibration factor 0.999 999 817, as reported in table 3 of Ref. [30].

Related research

People also read | Recommended articles | Cited by 30

The v3 band of 14C16O2 molecule measured by optical-frequency-comb-assisted cavity ring-down spectroscopy

Iacopo Galli et al.

Molecular Physics
Published online: 17 Oct 2011

Information for

Authors
R&D professionals
Editors
Librarians
Societies

Open access

Overview
Open journals
Open Select
Dove Medical Press
F1000Research

Opportunities

Reprints and e-prints
Advertising solutions
Accelerated publication
Corporate access solutions

Help and information

Help and contact
Newsroom
All journals
Books

Keep up to date

Register to receive personalised research and resources by email

Sign me up

Copyright © 2026 Informa UK Limited Privacy policy

Cookies Terms & conditions Accessibility

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG