



Molecular Physics >

An International Journal at the Interface Between Chemistry and Physics


Volume 117, 2019 - [Issue 3](#)193 | 21 | 0
Views | CrossRef citations to date | Altmetric

Research Article

Competition between σ -hole pnictogen bond and π -hole tetrel bond in complexes of $\text{CF}_2=\text{CFZH}_2$ (Z = P, As, and Sb)

Wenbo Dong, Yu Wang, Jianbo Cheng, Xin Yang & Qingzhong Li

Pages 251-259 | Received 12 Jun 2018, Accepted 24 Jul 2018, Published online: 12 Aug 2018

 Cite this article <https://doi.org/10.1080/00268976.2018.1508782>Sample our
Physical Sciences
Journals>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

Full Article

Figures & data

References

Supplemental

Citations

Metrics

Reprints & Permissions

Read this article

Share

ABSTRACT


A computational study of the complexes formed by $\text{F}_2\text{C}=\text{CFZH}_2$ (Z = P, As, and Sb) and $\text{F}_2\text{C}=\text{CFPF}_2$ with two Lewis bases (NH_3 and NMe_3) has been carried out. In general, two minima complexes are found, one with a σ -hole pnictogen bond and the other one with a π -hole tetrel bond in most complexes but two σ -hole pnictogen bonded complexes are obtained for $\text{F}_2\text{C}=\text{CFZH}_2$ and NH_3 . They have similar stability though $\text{F}_2\text{C}=\text{CFSbH}_2$ engages in a much stronger σ -hole pnictogen bond with NMe_3 . The $-\text{PF}_2$ substitution makes the π -hole on the terminal carbon form a tetrel bond with NH_3 . A heavier $-\text{ZH}_2$ group engages in a stronger σ -hole pnictogen bond but results in a weaker π -hole tetrel bond. Other than electrostatic interaction, the stability of both complexes is attributed

to the charge transfer from the N lone pair into the C-Z/H-Z anti-bonding orbital in the pnictogen bond and the C=C anti-bonding orbital in the tetrel bond.

The σ -hole pnictogen bonded and π -hole tetrel bonded complexes between $F_2C=CFZH_2$ (Z = P, As, and Sb) and two Lewis bases (NH_3 and NMe_3) have been compared. The results indicate that both interactions can compete, dependent on the nature of the N base.

GRAPHICAL ABSTRACT

σ -hole pnictogen bond



π -hole tetrel bond

KEYWORDS:

σ -hole pnictogen bond

π -hole tetrel bond

NBO

AIM

Disclosure statement

No potential conflict of interest was reported by the authors.

Additional information

Funding

This work was supported by National Natural Science Foundation of China [21573188].

Related Research Data

Computational insights into structural and optical properties of P-containing and N-containing ligands capped CdSe clusters

Source: Molecular Simulation

Binding Mechanisms in Supramolecular Complexes

Source: Angewandte Chemie International Edition

Characterizing the interplay of Pauli repulsion, electrostatics, dispersion and charge transfer in halogen bonding with energy decomposition analysis

Source: Physical Chemistry Chemical Physics

VMD: Visual molecular dynamics

Source: Journal of Molecular Graphics

Competition and cooperativity between tetrel bond and chalcogen bond in complexes involving F₂CX (X = Se and Te)

Source: Chemical Physics Letters

Comparison of σ -Hole and π -Hole Tetrel Bonds Formed by Pyrazine and 1,4-Dicyanobenzene: The Interplay between Anion- π and Tetrel Bonds



Related research 

People also read

Recommended articles

Cited by
21

Information for

Authors
R&D professionals
Editors
Librarians
Societies

Opportunities

Reprints and e-prints
Advertising solutions
Accelerated publication
Corporate access solutions

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG

Open access

Overview
Open journals
Open Select
Dove Medical Press
F1000Research

Help and information

Help and contact
Newsroom
All journals
Books

 Taylor and Francis
Group