









- Journal of Statistical Computation and Simulation
 List of Issues
 Volume 78, Issue 6
- Sampling nested Archimedean copulas

Journal of Statistical Computation and Simulation >

Volume 78, 2008 - <u>Issue 6</u>

981 | 175 | 0

Views CrossRef citations to date Altmetric

Original Articles

Sampling nested Archimedean copulas

Alexander J. McNeil

Pages 567-581 | Received 18 Jul 2006, Published online: 21 May 2008



Abstract

Full Article

Reprints & Permissions

We give algorithms for sampling from non-exchangeable Archimedean copulas created by the nesting of Archimedean copula generators, where in the most general algorithm the generators may be nested to an arbitrary depth. These algorithms are based on mixture representations of these copulas using Laplace transforms. While in principle the approach applies to all nested Archimedean copulas, in practice the approach is restricted to certain cases where we are able to sample distributions with given Laplace transforms. Precise instructions are given for the case when all generators are taken from the Gumbel parametric family or the Clayton family; the Gumbel case in particular proves very easy to simulate.

Keywords:

Archimedean copulas

Laplace transforms

Stochastic simulation

00A72 11K45 60E10 62E15 65C05

Acknowledgements

AMS 2000 MSC:

I thank Christoph Hummel of Converium for motivating this problem in the context of modelling dependent insurance losses. My thanks also go to Martin Pfister and Johanna Nešlehová for many useful discussions.

Additional information

Notes on contributors

Alexander J. McNeil

*Email: A.J.McNeil@hw.ac.uk

Related Research Data

Sampling from Archimedean copulas

Source: Quantitative Finance

An Introduction to Copulas

Source: Unknown Repository

Copula Methods in Finance

Source: Unknown Repository

Algorithms for Generating Discrete Random Variables with a Given Generating

Function or a Given Moment Sequence

Source: SIAM Journal on Scientific and Statistical Computing

Families of Multivariate Distributions

Source: Journal of the American Statistical Association A probabilistic interpretation of complete monotonicity

Source: Aequationes Mathematicae

On random variate generation when only moments or Fourier coefficients are known

Related research 1

People also read

Recommended articles

Cited by 175

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up











Accessibility



Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions



Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG