

1,021 Views | 179 CrossRef citations to date | 0 Altmetric

Original Articles

Sampling nested Archimedean copulas

Alexander J. McNeil

Pages 567-581 | Received 18 Jul 2006, Published online: 21 May 2008

🗨️ Cite this article 🔗 <https://doi.org/10.1080/00949650701255834>

Sample our
Computer Science
Journals

>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

📄 Full Article 📊 Figures & data 📖 References 🗨️ Citations 📊 Metrics

📄 Reprints & Permissions

Read this article

🔗 Share

Abstract

We give algorithms for sampling from non-exchangeable Archimedean copulas created by the nesting of Archimedean copula generators, where in the most general algorithm the generators may be nested to an arbitrary depth. These algorithms are based on mixture representations of these copulas using Laplace transforms. While in principle the approach applies to all nested Archimedean copulas, in practice the approach is restricted to certain cases where we are able to sample distributions with given Laplace transforms. Precise instructions are given for the case when all generators are taken from the Gumbel parametric family or the Clayton family; the Gumbel case in particular proves very easy to simulate.

Keywords:

Archimedean copulas

Laplace transforms

Stochastic simulation

00A72

11K45

60E10

62E15

65C05

Acknowledgements

I thank Christoph Hummel of Converium for motivating this problem in the context of modelling dependent insurance losses. My thanks also go to Martin Pfister and Johanna Nešlehová for many useful discussions.

Additional information

Notes on contributors

Alexander J. McNeil

*Email: A.J.McNeil@hw.ac.uk

Related Research Data

[Sampling from Archimedean copulas](#)

Source: Quantitative Finance

[An Introduction to Copulas](#)

Source: Unknown Repository

[Copula Methods in Finance](#)

Source: Unknown Repository

[Algorithms for Generating Discrete Random Variables with a Given Generating Function or a Given Moment Sequence](#)

Source: SIAM Journal on Scientific and Statistical Computing

[Families of Multivariate Distributions](#)

Source: Journal of the American Statistical Association

[A probabilistic interpretation of complete monotonicity](#)

Source: Aequationes Mathematicae

[On random variate generation when only moments or Fourier coefficients are known](#)

Related research

People also read

Recommended articles

Cited by
179

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG



Taylor & Francis
by informa