Journal of Statistical Computation and Simulation ›

Volume 78, 2008 - Issue 6

1,039 Views | 183 CrossRef citations to date | 0 Altmetric

Original Articles

Sampling nested Archimedean copulas

Alexander J. McNeil

Pages 567-581 | Received 18 Jul 2006, Published online: 21 May 2008

Cite this article https://doi.org/10.1080/00949650701255834

Sample our Computer Science Journals

>> Sign in here to start your access to the latest two volumes for 14 days

Abstract

We give algorithms for sampling from non-exchangeable Archimedean copulas created by the nesting of Archimedean copula generators, where in the most general algorithm the generators may be nested to an arbitrary depth. These algorithms are based on mixture representations of these copulas using Laplace transforms. While in principle the approach applies to all nested Archimedean copulas, in practice the approach is restricted to certain cases where we are able to sample distributions with given Laplace transforms. Precise instructions are given for the case when all generators are taken from the Gumbel parametric family or the Clayton family; the Gumbel case in particular proves very easy to simulate.

Keywords:

Archimedean copulas | Laplace transforms | Stochastic simulation

AMS 2000 MSC:

00A72 11K45 60E10 62E15 65C05

Previous article View issue table of contents **Next** article

Acknowledgements

I thank Christoph Hummel of Converium for motivating this problem in the context of modelling dependent insurance losses. My thanks also go to Martin Pfister and Johanna Nešlehová for many useful discussions.

Additional information

Notes on contributors

Alexander J. McNeil
*Email: A.J.McNeil@hw.ac.uk

Related Research Data

Sampling from Archimedean copulas
Source: Quantitative Finance
An Introduction to Copulas
Source: Unknown Repository
Copula Methods in Finance
Source: Unknown Repository
Algorithms for Generating Discrete Random Variables with a Given Generating Function or a Given Moment Sequence
Source: SIAM Journal on Scientific and Statistical Computing
Families of Multivariate Distributions
Source: Journal of the American Statistical Association
A probabilistic interpretation of complete monotonicity
Source: Aequationes Mathematicae
On random variate generation when only moments or Fourier coefficients are known

Source: Mathematics and Computers in Simulation

Related research

People also read | Recommended articles | Cited by 182

Hierarchies of Archimedean copulas

Cornelia Savu et al.
Quantitative Finance
Published online: 23 Jul 2009

A stochastic representation and sampling algorithm for nested Archimedean copulas

Marius Hofert
Journal of Statistical Computation and Simulation
Published online: 27 Jun 2011

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

Sign me up

Copyright © 2026 Informa UK Limited Privacy policy

Cookies Terms & conditions Accessibility

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG