



1,015 177

Views CrossRef citations to date Altmetric

0

Original Articles

Sampling nested Archimedean copulas

Alexander J. McNeil

Pages 567-581 | Received 18 Jul 2006, Published online: 21 May 2008

🗨️ Cite this article 🔗 <https://doi.org/10.1080/00949650701255834>

Sample our
Mathematics & Statistics
Journals



>> **Sign in here** to start your access
to the latest two volumes for 14 days

📄 Full Article

🖼️ Figures & data

📖 References

🗨️ Citations

📊 Metrics

🖨️ Reprints & Permissions

Read this article

🔗 Share

Abstract

We give algorithms for sampling from non-exchangeable Archimedean copulas created by the nesting of Archimedean copula generators, where in the most general algorithm the generators may be nested to an arbitrary depth. These algorithms are based on mixture representations of these copulas using Laplace transforms. While in principle the approach applies to all nested Archimedean copulas, in practice the approach is restricted to certain cases where we are able to sample distributions with given Laplace transforms. Precise instructions are given for the case when all generators are taken from the Gumbel parametric family or the Clayton family; the Gumbel case in particular proves very easy to simulate.

Keywords:

Archimedean copulas

Laplace transforms

Stochastic simulation

Acknowledgements

I thank Christoph Hummel of Converium for motivating this problem in the context of modelling dependent insurance losses. My thanks also go to Martin Pfister and Johanna Nešlehová for many useful discussions.

Additional information

Notes on contributors

Alexander J. McNeil
*Email: A.J.McNeil@hw.ac.uk

Related Research Data

[Sampling from Archimedean copulas](#)

Source: Quantitative Finance

[An Introduction to Copulas](#)

Source: Unknown Repository

[Copula Methods in Finance](#)

Source: Unknown Repository

[Algorithms for Generating Discrete Random Variables with a Given Generating Function or a Given Moment Sequence](#)

Source: SIAM Journal on Scientific and Statistical Computing

[Families of Multivariate Distributions](#)

Source: Journal of the American Statistical Association

[A probabilistic interpretation of complete monotonicity](#)

Source: Aequationes Mathematicae

[On random variate generation when only moments or Fourier coefficients are known](#)

Related research 

People also read

Recommended articles

Cited by
177

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up

