

▶ Volume 88, Issue 12 ▶ Variance Inflation Factor and Condition ....

Journal of Statistical Computation and Simulation > Volume 88, 2018 - Issue 12

4.251 139

Views CrossRef citations to date Altmetric

**Original Articles** 

# Variance Inflation Factor and Condition Number in multiple linear regression

R. Salmerón 📵, C. B. García 🔀 📵 & J. García

Pages 2365-2384 | Received 19 Apr 2017, Accepted 07 Apr 2018, Published online: 22 Apr 2018

▶ https://doi.org/10.1080/00949655.2018.1463376 **66** Cite this article



Sample our Computer Science

Full Article

Figures & data

References

**66** Citations

**Metrics** 

Reprints & Permissions

Read this article

## ABSTRACT

The Variance Inflation Factor and the Condition Number are measures traditionally applied to detect the presence of collinearity in a multiple linear model. This paper presents the relation and the difference between both measures from theoretical and empirical perspectives by using Monte Carlo simulations and taking special interest in the computational techniques.

#### **Q** KEYWO

#### About Cookies On This Site

We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our Privacy Policy

Accept All **Essential Onl** Settings

No poter




## **ORCID**

- R. Salmerón <a href="http://orcid.org/0000-0003-2589-4058">http://orcid.org/0000-0003-2589-4058</a>
- C. B. García <a href="http://orcid.org/0000-0003-1622-3877">http://orcid.org/0000-0003-1622-3877</a>

## Notes

one.

- 1 Note that the constant term disappears after the standardization of the data.
- 2 Note that, when data are standardized, the VIF and CN coincide with the result obtained from typified data.
- 3 Note that these examples are not regression models since n=p.
- 4 Denoting X1=1, the auxiliary regression to calculate the VIF is expressed as  $X2=\gamma1+w$ , where it is verified that  $\gamma^*=X^*-2$  and, consequently,  $SSR=\Sigma i=1n(X2i-X^*-2)2=SST$ . In this case, it is always verified that Raux2=1. The version of the previous regression with unit length data is given by X2,  $Iu=\gamma1Iu+w$  where X2, Iu=X/a with  $a=\Sigma i=1nX2i2$  and Iu=1/n. In this case,  $\gamma^*=n/a\cdot X^*-2$  and, then,  $SSR=(1/a)\Sigma i=1n(X2i-n\cdot X^*-2\cdot 1/n)2=SST$ . Thus, this situation will be similar to the initial



Information for

Authors

**R&D** professionals

**Editors** 

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

**Open Select** 

**Dove Medical Press** 

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

## Keep up to date

Register to receive personalised research and resources by email



Sign me up











Copyright © 2024 Informa UK Limited Privacy policy Cookies Terms & conditions



Accessibility

Registered in England & Wales No. 3099067 5 Howick Place | London | SW1P 1WG

### About Cookies On This Site



We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our <a href="Privacy Policy">Privacy Policy</a>



Essential Onl

Settings