

5,079 Views | 267 CrossRef citations to date | 2 Altmetric

Original Articles

Variance Inflation Factor and Condition Number in multiple linear regression

R. Salmerón , C. B. García   & J. García

Pages 2365-2384 | Received 19 Apr 2017, Accepted 07 Apr 2018, Published online: 22 Apr 2018

 Cite this article  <https://doi.org/10.1080/00949655.2018.1463376>



Sample our
Computer Science
Journals

>> **Sign in here** to start your access
to the latest two volumes for 14 days

 Full Article  Figures & data  References  Citations  Metrics

 Reprints & Permissions

Read this article

Share

ABSTRACT

The Variance Inflation Factor and the Condition Number are measures traditionally applied to detect the presence of collinearity in a multiple linear model. This paper presents the relation and the difference between both measures from theoretical and empirical perspectives by using Monte Carlo simulations and taking special interest in the computational techniques.

KEYWORDS:

Variance Inflation Factor

Condition Number

multicollinearity detection

data transformation

 Previous article

View issue table of contents

Next article 

Disclosure statement

No potential conflict of interest was reported by the authors.

ORCID

R. Salmerón <http://orcid.org/0000-0003-2589-4058>

C. B. García <http://orcid.org/0000-0003-1622-3877>

Notes

- 1 Note that the constant term disappears after the standardization of the data.
- 2 Note that, when data are standardized, the VIF and CN coincide with the result obtained from typified data.
- 3 Note that these examples are not regression models since $n=p$.
- 4 Denoting $X_1=1$, the auxiliary regression to calculate the VIF is expressed as $X_2=\gamma_1+w$, where it is verified that $\hat{\gamma}=X^{-2}$ and, consequently, $SSR=\sum_{i=1}^n(X_{2i}-X^{-2})^2=SST$. In this case, it is always verified that $R_{aux}^2=1$. The version of the previous regression with unit length data is given by $X_{2,lu}=\gamma_1lu+w$ where $X_{2,lu}=X/a$ with $a=\sum_{i=1}^nX_{2i}^2$ and $1lu=1/n$. In this case, $\hat{\gamma}=n/a\cdot X^{-2}$ and, then, $SSR=(1/a)\sum_{i=1}^n(X_{2i}-n\cdot X^{-2}\cdot 1/n)^2=SST$. Thus, this situation will be similar to the initial one.

Related research

People also read

Recommended articles

Cited by
267

[Model-Dependent Variance Inflation Factor Cutoff Values >](#)

Trevor A. Craney et al.
Quality Engineering
Published online: 15 Feb 2007

Christopher Glen Thompson et al.
Basic and Applied Social Psychology
Published online: 1 Feb 2017

Information for

Authors
R&D professionals
Editors
Librarians
Societies

Opportunities

Reprints and e-prints
Advertising solutions
Accelerated publication
Corporate access solutions

Open access

Overview
Open journals
Open Select
Dove Medical Press
F1000Research

Help and information

Help and contact
Newsroom
All journals
Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG



Taylor & Francis
by informa