



Phase Transitions >

A Multinational Journal

Volume 81, 2008 - [Issue 5](#)

120 Views | 16 CrossRef citations to date | 0 Altmetric

Original Articles

Soft mode dispersion and 'waterfall' phenomenon in relaxors revisited

J. Hlinka  & M. Kempa

Pages 491-508 | Received 03 Dec 2007, Accepted 04 Feb 2008, Published online: 07 May 2008

 Cite this article

Sample our
Earth Sciences
Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

 Full Article

 Figures & data

 References

 Citations

 Metrics

 Reprints & Permissions

Read this article

 Share

Abstract

Results of recent inelastic neutron scattering studies of lead-based relaxor ferroelectrics by Gvasaliya et al. [J. Phys.: Condens. Matter 17, 4343 (2005); J. Phys.: Condens. Matter 19, 016219 (2007)] have put in question the existence of the “waterfall” anomaly—an apparent vertical dispersion segment joining the TA and TO branches—observed earlier in low-energy $[\xi 00]$ phonon dispersion curves of these materials. In the present article, we review the results of earlier experiments and model calculations together with the outcome of our recent measurements on PMN using the same instrumental set-up as Gvasaliya et al. to conclude that the “waterfall” feature is not an experimental artefact. We also give some hints on a possible explanation of the results of Gvasaliya et al., by exploring the fact that the reported dispersion of the underdamped transverse optic branch follows the longitudinal acoustic (LA) branch dispersion surprisingly closely.

Keywords:

Acknowledgements

The work is partly based on experiments performed at the Swiss spallation neutron source SINQ, Paul Scherrer Institute, Villigen, Switzerland. This research project has been supported by the European Commission under the 6th Framework Programme through the Key Action: Strengthening the European Research Area, Research Infrastructures. Contract No.: RII3-CT-2003-505925. Additional financial support was obtained from the Czech grant agencies (projects GACR 202/06/0403 and GACR 202/06/0411 and AVOZ 10100520). Authors thank to Dr S. Gvasaliya from Laboratory for Neutron Scattering of ETH Zurich and Paul Scherrer Institut in Villigen for useful discussions and the help with set-up of the experiment at TASP instrument, to Dr J. Petzelt from the Institute of Physics of Czech Academy of Science in Prague for careful reading of the manuscript and to Dr J. Kulda from the Institute Laue-Langevin for his long-term interest, encouragement, valuable suggestions and numerous critical discussions.

Notes

Notes

1. In the rest of this article, we use reduced (dimensionless) phonon wave vectors expressed in reciprocal lattice units (r.l.u.) given by $c^* = 2\pi/c$, (e.g. for PMN, $c = 4.04 \text{ \AA}$, i.e. $c^* = 2\pi/c = 1.555 \text{ \AA}^{-1}$).
2. First two hypotheses are close to those invoked in Ref. [17](#).
3. Apart from the obvious misprint-extra factor of 2 in arguments of all trigonometric functions introduced in Ref. [14](#).
4. This additional mode should not be confused with additional relaxations in the GHz region, often called (narrow) central peaks, which are not in the scope of this article.

5. We have combined the data from the (200) and (300) BZs as D_q should be independent of the BZ choice.
6. In principle, PMN is known to grow also, for example in pyrochlore structure, but in this case both the lattice parameters and the TO mode frequency are completely different.
7. It was argued in [9](#) that the TO mode cannot couple noticeably to the TA branch because the independent mode intensities do not change with temperature. However, this not a valid argument since the measurements shown in figure of Ref. [9](#) were done in (20q) zone, where both TA and TO modes have similar structure factors so that eventual eigenvector change has no chance to produce such drastic intensity changes as those observed in the quoted [9](#) case of SrTiO_3 .

Related Research Data

[The rhombohedral phase with incommensurate modulation in \$\text{Na}_{1/2}\text{Bi}_{1/2}\text{TiO}_3\$](#)

Source: Phase Transitions

[Broad-band dielectric response of \$\text{PbMg}_{1/3}\text{Nb}_{2/3}\text{O}_3\$ relaxor ferroelectrics: Single crystals, ceramics and thin films](#)

Source: Journal of the European Ceramic Society

[Neutron scattering studies of the mechanism of ferroelectricity in \$68\%\text{PbMg}_{1/3}\text{Nb}_{2/3}\text{O}_3\$ - \$32\%\text{PbTiO}_3\$](#)

Source: Journal of Physics Condensed Matter

[Origin of the “Waterfall” Effect in Phonon Dispersion of Relaxor Perovskites](#)

Source: Physical Review Letters

[Distribution of correlation radii in disordered ferroelectrics](#)

Source: Applied Physics Letters

[Inelastic Neutron Scattering from Single-Domain \$\text{BaTiO}_3\$](#)

Source: Unknown Repository

[The Lineshape of Inelastic Neutron Scattering in Relaxor Ferroelectrics](#)

Source: Physics of the Solid State

[Soft and central mode behaviour in \$\text{PbMg}_{1/3}\text{Nb}_{2/3}\text{O}_3\$ relaxor ferroelectric](#)

Source: Journal of Physics Condensed Matter

[Mode coupling and polar nanoregions in the relaxor ferroelectric \$\text{Pb}\(\text{Mg}_{1/3}\text{Nb}_{2/3}\)\text{O}_3\$](#)

Source: Unknown Repository

[Soft Phonon Anomalies in the Relaxor Ferroelectric \$\text{Pb}\(\text{Zn}_{1/3}\text{Nb}_{2/3}\)_{0.92}\text{Ti}_{0.08}\text{O}_3\$](#)

Source: Physical Review Letters

Soft Mode Dynamics above and below the Burns Temperature in the Relaxor $\text{Pb}(\text{Mg}_{1/3}\text{Nb}_{2/3})\text{O}_3$

Source: Physical Review Letters

Direct evidence of soft mode behavior near the Burns temperature in the $\text{PbMg}_{1/3}\text{Nb}_{2/3}\text{O}_3$ relaxor ferroelectric

Source: Unknown Repository

Phonons and fractons in the vibration spectrum of the relaxor ferroelectric $\text{PbMg}_{1/3}\text{Nb}_{2/3}\text{O}_3$

Source: Physica B Condensed Matter

Dirty displacive ferroelectrics

Source: Unknown Repository

Diffuse scattering in $\text{Pb}(\text{Zn}_{1/3}\text{Nb}_{2/3})\text{O}_3$ with 8 PbTiO_3 by quasi-elastic neutron scattering

Source: Journal of Physics Condensed Matter

Strong Influence of the Diffuse Component on the Lattice Dynamics in $\text{Pb}(\text{Mg}_{1/3}\text{Nb}_{2/3})\text{O}_3$

Source: Journal of the Physical Society of Japan

Neutron Scattering with a Triple-Axis Spectrometer

Source: Unknown Repository

Quasi-elastic scattering, random fields and phonon-coupling effects in $\text{PbMg}_{1/3}\text{Nb}_{2/3}\text{O}_3$

Source: Journal of Physics Condensed Matter

Infrared and Raman spectroscopy of $[\text{Pb}(\text{Zn}_{1/3}\text{Nb}_{2/3})\text{O}_3]_{0.92}-[\text{PbTiO}_3]_{0.08}$ and $[\text{Pb}(\text{Mg}_{1/3}\text{Nb}_{2/3})\text{O}_3]_{0.71}-[\text{PbTiO}_3]_{0.29}$ single crystals

Source: Journal of Applied Physics

Polar modes in relaxor $\text{PbMg}_{1/3}\text{Nb}_{2/3}\text{O}_3$ by hyper-Raman scattering

Source: Physical Review B

Inelastic neutron scattering study of the relaxor ferroelectric PbMg Nb O at high temperatures

Source: The European Physical Journal B

Dynamical effects of the nanometer-sized polarized domains in $\text{Pb}(\text{Zn}_{1/3}\text{Nb}_{2/3})\text{O}_3$

Source: Unknown Repository

Crystal Dynamics of Lead. I. Dispersion Curves at 100°K

Source: Unknown Repository

Soft mode anomalies in the perovskite relaxor $\text{Pb}(\text{Mg}_{1/3}\text{Nb}_{2/3})\text{O}_3$

Source: Unknown Repository

Raman scattering from in-plane lattice modes in low-stage graphite-alkali-metal compounds

Source: Unknown Repository

Anisotropic Dielectric Function in Polar Nanoregions of Relaxor Ferroelectrics

Source: Physical Review Letters

Phase diagram of the relaxor ferroelectric $(1-x)\text{Pb}(\text{Zn}_{1/3}\text{Nb}_{2/3})\text{O}_3-x\text{PbTiO}_3$

Source: Unknown Repository

Hyper-Raman-active soft mode in $\text{Pb}(\text{Mg}_{1/3}\text{Nb}_{2/3})_{0.73}\text{Ti}_{0.27}\text{O}_3$

Source: Physical Review B

Lattice Dynamics of Disordered Perovskite $\text{Pb}(\text{Zn}_{1/3}\text{Nb}_{2/3})\text{O}_3$

Source: Journal of the Physical Society of Japan

Phonons in $\text{PbMg}_{1/3}\text{Nb}_{2/3}\text{O}_3$ Measured by Inelastic Neutron Scattering

Source: Ferroelectrics

Disorder and relaxation mode in the lattice dynamics of the $\text{PbMg}_{1/3}\text{Nb}_{2/3}\text{O}_3$ relaxor ferroelectric

Source: Physical Review B

Anomalous transverse acoustic phonon broadening in the relaxor ferroelectric $\text{Pb}(\text{Mg}_{1/3}\text{Nb}_{2/3})_{0.8}\text{Ti}_{0.2}\text{O}_3$

Source: Unknown Repository

Soft-mode spectroscopy: Experimental studies of structural phase transitions

Source: Reviews of Modern Physics

Nonferroelectric aging in the relaxor $\text{PbMg}_{1/3}\text{Nb}_{2/3}\text{O}_3$

Source: Physical Review B

Linking provided by 

Related research

People also read

Recommended articles

Cited by
16

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



[Sign me up](#)



[Copyright © 2025](#) [Informa UK Limited](#) [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)



Taylor & Francis Group
an **informa** business

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG