



Phase Transitions >

A Multinational Journal

Volume 81, 2008 - [Issue 5](#)

132 | 16

Views | CrossRef citations to date | 0 | Altmetric

Original Articles

# Soft mode dispersion and 'waterfall' phenomenon in relaxors revisited

J. Hlinka  & M. Kempa

Pages 491-508 | Received 03 Dec 2007, Accepted 04 Feb 2008, Published online: 07 May 2008

 Cite this article

Sample our  
Earth Sciences  
Journals

>> [Sign in here](#) to start your access  
to the latest two volumes for 14 days



 Full Article

 Figures & data

 References

 Citations

 Metrics

 Reprints & Permissions

Read this article

 Share

## Abstract

Results of recent inelastic neutron scattering studies of lead-based relaxor ferroelectrics by Gvasaliya et al. [J. Phys.: Condens. Matter 17, 4343 (2005); J. Phys.: Condens. Matter 19, 016219 (2007)] have put in question the existence of the “waterfall” anomaly—an apparent vertical dispersion segment joining the TA and TO branches—observed earlier in low-energy  $[\xi 00]$  phonon dispersion curves of these materials. In the present article, we review the results of earlier experiments and model calculations together with the outcome of our recent measurements on PMN using the same instrumental set-up as Gvasaliya et al. to conclude that the “waterfall” feature is not an experimental artefact. We also give some hints on a possible explanation of the results of Gvasaliya et al., by exploring the fact that the reported dispersion of the underdamped transverse optic branch follows the longitudinal acoustic (LA) branch dispersion surprisingly closely.

relaxor ferroelectrics

waterfall phenomenon

phonon dispersions

inelastic neutron scattering

---

## Acknowledgements

The work is partly based on experiments performed at the Swiss spallation neutron source SINQ, Paul Scherrer Institute, Villigen, Switzerland. This research project has been supported by the European Commission under the 6th Framework Programme through the Key Action: Strengthening the European Research Area, Research Infrastructures. Contract No.: RII3-CT-2003-505925. Additional financial support was obtained from the Czech grant agencies (projects GACR 202/06/0403 and GACR 202/06/0411 and AVOZ 10100520). Authors thank to Dr S. Gvasaliya from Laboratory for Neutron Scattering of ETH Zurich and Paul Scherrer Institut in Villigen for useful discussions and the help with set-up of the experiment at TASP instrument, to Dr J. Petzelt from the Institute of Physics of Czech Academy of Science in Prague for careful reading of the manuscript and to Dr J. Kulda from the Institute Laue-Langevin for his long-term interest, encouragement, valuable suggestions and numerous critical discussions.

---

## Notes

### Notes

1. In the rest of this article, we use reduced (dimensionless) phonon wave vectors expressed in reciprocal lattice units (r.l.u.) given by  $c^* = 2\pi/c$ , (e.g. for PMN,  $c = 4.04 \text{ \AA}$ , i.e.  $c^* = 2\pi/c = 1.555 \text{ \AA}^{-1}$ ).
2. First two hypotheses are close to those invoked in Ref. [17](#).
3. Apart from the obvious misprint-extra factor of 2 in arguments of all trigonometric functions introduced in Ref. [14](#).
4. This additional mode should not be confused with additional relaxations in the GHz region, often called (narrow) central peaks, which are not in the scope of this article.

5. We have combined the data from the (200) and (300) BZs as  $D_q$  should be independent of the BZ choice.
6. In principle, PMN is known to grow also, for example in pyrochlore structure, but in this case both the lattice parameters and the TO mode frequency are completely different.
7. It was argued in [9](#) that the TO mode cannot couple noticeably to the TA branch because the independent mode intensities do not change with temperature. However, this not a valid argument since the measurements shown in figure of Ref. [9](#) were done in (20q) zone, where both TA and TO modes have similar structure factors so that eventual eigenvector change has no chance to produce such drastic intensity changes as those observed in the quoted [9](#) case of  $\text{SrTiO}_3$ .

#### Related Research Data

[The rhombohedral phase with incommensurate modulation in  \$\text{Na}\_{1/2}\text{Bi}\_{1/2}\text{TiO}\_3\$](#)

Source: Phase Transitions

[Broad-band dielectric response of  \$\text{PbMg}\_{1/3}\text{Nb}\_{2/3}\text{O}\_3\$  relaxor ferroelectrics: Single crystals, ceramics and thin films](#)

Source: Journal of the European Ceramic Society

[Neutron scattering studies of the mechanism of ferroelectricity in  \$68\%\text{PbMg}\_{1/3}\text{Nb}\_{2/3}\text{O}\_3\$ - \$32\%\text{PbTiO}\_3\$](#)

Source: Journal of Physics Condensed Matter

[Origin of the “Waterfall” Effect in Phonon Dispersion of Relaxor Perovskites](#)

Source: Physical Review Letters

[Distribution of correlation radii in disordered ferroelectrics](#)

Source: Applied Physics Letters

[Inelastic Neutron Scattering from Single-Domain  \$\text{BaTiO}\_3\$](#)

Source: Unknown Repository

[The Lineshape of Inelastic Neutron Scattering in Relaxor Ferroelectrics](#)

#### Related research

People also read

Recommended articles

Cited by  
16

## Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

## Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

## Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

## Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

## Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954  
5 Howick Place | London | SW1P 1WG



**Taylor & Francis**  
by informa