







Q

Home ► All Journals ► Physical Sciences ► Phase Transitions ► List of Issues ► Volume 81, Issue 5 ► Soft mode dispersion and 'waterfall' phe

Phase Transitions >

A Multinational Journal

Volume 81, 2008 - <u>Issue 5</u>

120 | 16 | 0

Views CrossRef citations to date Altmetric

Original Articles

Soft mode dispersion and 'waterfall' phenomenon in relaxors revisited

I. Hlinka 🔀 & M. Kempa

Pages 491-508 | Received 03 Dec 2007, Accepted 04 Feb 2008, Published online: 07 May 2008

66 Cite this article



Abstract

Results of recent inelastic neutron scattering studies of lead-based relaxor ferroelectrics by Gvasaliya et al. [J. Phys.: Condens. Matter 17, 4343 (2005); J. Phys.: Condens. Matter 19, 016219 (2007)] have put in question the existence of the "waterfall" anomaly-an apparent vertical dispersion segment joining the TA and TO branches-observed earlier in low-energy [§00] phonon dispersion curves of these materials. In the present article, we review the results of earlier experiments and model calculations together with the outcome of our recent measurements on PMN using the same instrumental set-up as Gvasaliya et al. to conclude that the "waterfall" feature is not an experimental artefact. We also give some hints on a possible explanation of the results of Gvasaliya et al., by exploring the fact that the reported dispersion of the underdamped transverse optic branch follows the longitudinal acoustic (LA) branch dispersion surprisingly closely.

Keywords:

relaxor ferroelectrics waterfall phenomenon phonon dispersions inelastic neutron scattering

Acknowledgements

The work is partly based on experiments performed at the Swiss spallation neutron source SINQ, Paul Scherrer Institute, Villigen, Switzerland. This research project has been supported by the European Commission under the 6th Framework Programme through the Key Action: Strengthening the European Research Area, Research Infrastructures. Contract No.: RII3-CT-2003-505925. Additional financial support was obtained from the Czech grant agencies (projects GACR 202/06/0403 and GACR 202/06/0411 and AVOZ 10100520). Authors thank to Dr S. Gvasaliya from Laboratory for Neutron Scattering of ETH Zurich and Paul Scherrer Institut in Villigen for useful discussions and the help with set-up of the experiment at TASP instrument, to Dr J. Petzelt from the Institute of Physics of Czech Academy of Science in Prague for careful reading of the manuscript and to Dr J. Kulda from the Institute Laue-Langevin for his long-term interest, encouragement, valuable suggestions and numerous critical discussions.

Notes

Notes

- 1. In the rest of this article, we use reduced (dimensionless) phonon wave vectors expressed in reciprocal lattice units (r.l.u.) given by $c^* = 2\pi/c$, (e.g. for PMN, c = 4.04 Å, i.e. $c^* = 2\pi/c = 1.555$ Å⁻¹).
- 2. First two hypotheses are close to those invoked in Ref. 17.
- 3. Apart from the obvious misprint–extra factor of 2 in arguments of all trignometric functions introduced in Ref. <u>14</u>.
- 4. This additional mode should not be confused with additional relaxations in the GHz region, often called (narrow) central peaks, which are not in the scope of this article.

- 5. We have combined the data from the (200) and (300) BZs as D $_{\rm q}$ should be independent of the BZ choice.
- 6. In principle, PMN is known to grow also, for example in pyrochlore structure, but in this case both the lattice parameters and the TO mode frequency are completely different.
- 7. It was argued in $\underline{9}$ that the TO mode cannot couple noticeably to the TA branch because the independent mode intensities do not change with temperature. However, this not a valid argument since the measurements shown in figure of Ref. $\underline{9}$ were done in (20q) zone, where both TA and TO modes have similar structure factors so that eventual eigenvector change has no chance to produce such drastic intensity changes as those observed in the quoted $\underline{9}$ case of SrTiO₃.

Related Research Data

The rhombohedral phase with incommensurate modulation in Na_{1/2}Bi_{1/2}TiO₃

Source: Phase Transitions

Broad-band dielectric response of PbMg1/3Nb2/3O3 relaxor ferroelectrics: Single

crystals, ceramics and thin films

Source: Journal of the European Ceramic Society

Neutron scattering studies of the mechanism of ferroelectricity in $68\%PbMg_{1/3}Nb_{2/3}O_3$ -

32%PbTiO₃

Source: Journal of Physics Condensed Matter

Origin of the "Waterfall" Effect in Phonon Dispersion of Relaxor Perovskites

Source: Physical Review Letters

Distribution of correlation radii in disordered ferroelectrics

Source: Applied Physics Letters

Inelastic Neutron Scattering from Single-Domain BaTiO3

Source: Unknown Repository

The Lineshape of Inelastic Neutron Scattering in Relaxor Ferroelectrics

Related research 1

People also read

Recommended articles

Cited by

16

R. A. Cowley et al.

Advances in Physics

Published online: 25 Mar 2011

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up











Accessibility



Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions

Taylor & Francis Group an informa business

Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG