

International Journal of Remote Sensing >

Volume 28, 2007 - [Issue 12](#)

630 Views | 52 CrossRef citations to date | 0 Altmetric

Original Articles

# Stabilizing high-order, non-classical harmonic analysis of NDVI data for average annual models by damping model roughness

J. F. Hermance 

Pages 2801-2819 | Received 05 May 2006, Accepted 06 Jul 2006, Published online: 29 May 2007

 Cite this article  <https://doi.org/10.1080/01431160600967128>

Sample our  
Geography  
Journals  
>> [Sign in here](#) to start your access  
to the latest two volumes for 14 days

 Full Article  Figures & data  References  Citations  Metrics

 Reprints & Permissions [Read this article](#) [Share](#)

## Abstract

Fourier series and related harmonic methods have been demonstrably effective for identifying and characterizing the seasonal behaviour, or phenology, of a variety of terrestrial vegetation communities using Normalized Difference Vegetation Index (NDVI) time series from Earth-orbiting satellites. The ultimate temporal resolution of such applications has been limited, however, by the common practice of truncating, or low pass filtering, harmonic series to relatively low order terms, in order to suppress spurious oscillations in the model results. The temporal resolution of these techniques can be significantly improved if, along with a weighted minimization of the sum of the squared data residuals tracking the upper envelope of observed data, we also enforce an expectation of minimum model roughness to dampen spurious oscillations in

predicted values. The resulting annual models have resolutions consistent with the application of special transcendental forms, such as asymmetric Gaussian and logistic (sigmoidal) functions, recently reported in the literature.

---

## Acknowledgments

This work was developed over several years of interacting and constructive discussions with Bethany Bradley and Rob Jacob, both recent recipients of PhD degrees from Brown University. Professor Jack Mustard generously supplied the AVHRR-NDVI data and the hospitable venue in which this work was possible. Financial support was provided by Brown University and private sector partners with its Environmental Geophysics/Hydrology group.

---

### Related Research Data

[Mapping agroecological zones and time lag in vegetation growth by means of fourier analysis of time series of NDVI images](#)

Source: *Advances in Space Research*

[Savitzky-Golay Smoothing Filters](#)

Source: *Computers in Physics*

[Improved monitoring of vegetation dynamics at very high latitudes: A new method using MODIS NDVI](#)

Source: *Remote Sensing of Environment*

[The 1 km AVHRR global land data set: first stages in implementation](#)

Source: *International Journal of Remote Sensing*

[A curve fitting procedure to derive inter-annual phenologies from time series of noisy satellite NDVI data](#)

Source: *Remote Sensing of Environment*

[Extracting Phenological Signals From Multiyear AVHRR NDVI Time Series: Framework for Applying High-Order Annual Splines With Roughness Damping](#)

People also read

Recommended articles

Cited by  
52

Spatio-Temporal Reconstruction of MODIS NDVI by Regional Land Surface Phenology and Harmonic Analysis of Time-Series >

Suman Kumar Padhee et al.  
GIScience & Remote Sensing  
Published online: 6 Aug 2019



## Information for

Authors

R&D professionals

Editors

Librarians

Societies

## Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

## Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

## Help and information

Help and contact

Newsroom

All journals

Books

## Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954  
5 Howick Place | London | SW1P 1WG



Taylor & Francis  
by informa