

Home ► All Journals ► Geography ► International Journal of Remote Sensing ► List of Issues ► Volume 28, Issue 12 ► Stabilizing high-order, non-classical ha ...

International Journal of Remote Sensing > Volume 28, 2007 - <u>Issue 12</u>

594500ViewsCrossRef citations to dateAltmetric

Original Articles

Stabilizing high-order, non-classical harmonic analysis of NDVI data for average annual models by damping model roughness

J. F. Hermance 🔽

Pages 2801-2819 | Received 05 May 2006, Accepted 06 Jul 2006, Published online: 29 May 2007

G Cite this article **I** https://doi.org/10.1080/01431160600967128

Sample our Environment & Agriculture journals, sign in here to start your access, latest two full volumes FREE to you for 14 days



Fourier s identifyi terrestria time ser applicati pass filspur can be s squared an expec predicte applicati (sigmoic We and our 909 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting I Accept enables tracking technologies to support the purposes shown under we and our partners process data to provide. Selecting Reject All or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the Show Purposes link on the bottom of the webpage .Your choices will have effect within our Website. For more details, refer to our Privacy Policy. <u>Here</u>

We and our partners process data to provide:

Use precise geolocation data. Actively scan device



Q

Acknowledgments

This work was developed over several years of interacting and constructive discussions with Bethany Bradley and Rob Jacob, both recent recipients of PhD degrees from Brown University. Professor Jack Mustard generously supplied the AVHRR-NDVI data and the hospitable venue in which this work was possible. Financial support was provided by Brown University and private sector partners with its Environmental Geophysics/Hydrology group.



Source: International Journal of Remote Sensing African Land-Cover Classification Using Satellite Data Source: Science Monitoring vegetation phenology using MODIS Source: Remote Sensing of Environment Seasonality extraction by function fitting to time-series of satellite sensor data Source: IEEE Transactions on Geoscience and Remote Sensing Fourier Series for analysis of temporal sequences of satellite sensor imagery Source: International Journal of Remote Sensing Land-Surface Phenologies from AVHRR Using the Discrete Fourier Transform Source: Remote Sensing of Environment Interannual variations in satellite-sensed vegetation index data from 1981 to 1991 Source: Journal of Geophysical Research Atmospheres Measuring phenological variability from satellite imagery Source: Journal of Vegetation Science Identifying land cover variability distinct from land cover change: Cheatgrass in the **Great Basin** Source: Remote Sensing of Environment A simple method for reconstructing a high-quality NDVI time-series data set based on the Savitzky-Golay filter Source: Remote Sensing of Environment Crop identification using harmonic analysis of time-series AVHRR NDVI data Sourc X TIMES Sourc ourier Mapp analy Sourc Gree :e Sourc obal Sour Class g Fouri Sourc tion Fouri Sourc

Related research (1)

People also read	Recommended articles	Cited by 50	
Information for	Open access		
Authors	Overview		
R&D professionals	Open journals	Open journals	
Editors	Open Select	Open Select	
Librarians	Dove Medical Press		
Societies	F1000Research	F1000Research	
Opportunities	Help and inform	Help and information	
Reprints and e-prints	Help and contact	Help and contact	
Advertising solutions	Newsroom	Newsroom	
Accelerated publication	All journals	All journals	
Corporate access solutions	Books		

Кеер ир



Accessib

Registered 5 Howick Pl

X

or & Francis Group orma business