

614 59 Views CrossRef citations to date 0 Altmetric

Original Articles

# Estimating above-ground biomass in young forests with airborne laser scanning

Erik Næsset

Pages 473-501 | Received 29 Apr 2009, Accepted 30 Sep 2009, Published online: 06 Feb 2011

Cite this article <https://doi.org/10.1080/01431160903474970>

Sample our Environment & Agriculture Journals

>> [Sign in here](#) to start your access to the latest two volumes for 14 days

- Full Article
- Figures & data
- References
- Citations
- Metrics
- Reprints & Permissions
- [Read this article](#)

## Abstract

Total above-ground biomass of spruce, pine and birch was estimated in three different field data sets. The biomass was estimated using airborne laser scanning (ALS) data with a point cloud (PC) derived from the ALS data. The biomass ranged from 200 to 232.9 t ha<sup>-1</sup> and the ALS-derived biomass ranged from 200 to 232.9 t ha<sup>-1</sup>. The ALS-derived biomass and ALS-derived biomass models were compared with the ALS-derived biomass models. The ALS-derived biomass models were compared with the ALS-derived biomass models. The ALS-derived biomass models were compared with the ALS-derived biomass models.

**We Care About Your Privacy**

We and our 843 partners store and/or access information on a device, such as unique IDs in cookies to process personal data. You may accept or manage your choices by clicking below, including your right to object where legitimate interest is used, or at any time in the privacy policy page. These choices will be signaled to our partners and will not affect browsing data. [Privacy Policy](#)

We and our partners process data to provide:

- Use precise geolocation data. Actively scan device characteristics for identification. Store and/or access information on a device. Personalised advertising and content, advertising and content measurement, audience research and services development.

List of Partners (vendors)

[I Accept](#)

[Essential Only](#)

[Show Purpose](#)



rights  
derived  
(S) data with  
200-232.9  
mass ranged  
) canopy  
and ALS-  
near models,  
variables, and  
by  
limit  
explained

variability of the estimated models ranged from 60% to 83%. Tree species had a significant influence on the models. For given values of the ALS-derived metrics related to canopy height and canopy density, spruce tended to have higher above-ground biomass values than pine and deciduous species. There were no clear effects of model form and canopy threshold on the accuracy of predictions produced by cross validation of the various models, but there is a risk of heteroskedasticity with linear models. Cross validation revealed an accuracy of the root mean square error (RMSE) ranging from 3.85 to 13.9 Mg ha<sup>-1</sup>, corresponding to 22.6% to 48.1% of mean field-measured biomass. It was concluded that airborne laser scanning has a potential for predicting biomass in young forest stands (> 0.5 ha) with an accuracy of 20–30% of mean ground value.

#### Related Research Data

[Airborne laser scanning as a method in operational forest inventory: Status of accuracy assessments accomplished in Scandinavia](#)

Source: Informa UK Limited

[Assessing Point Accuracy of DGPS Under Forest Canopy Before Data Acquisition, in the Field and After Postprocessing](#)

Source: Informa UK Limited

[Estimating tree heights and number of stems in young forest stands using airborne laser](#)

Source

Estim

latitu

predi

Source

A

St

Biom

Source

Estim

zone

Source



direct

rest

Estimation of above ground forest biomass from airborne discrete return laser scanner data using canopy-based quantile estimators

Source: Informa UK Limited

A Heteroskedasticity-Consistent Covariance Matrix Estimator and a Direct Test for Heteroskedasticity

Source: JSTOR

Surface lidar remote sensing of basal area and biomass in deciduous forests of eastern Maryland, USA

Source: Elsevier BV

Use of Large-Footprint Scanning Airborne Lidar To Estimate Forest Stand Characteristics in the Western Cascades of Oregon

Source: Elsevier BV

Effects of different sensors, flying altitudes, and pulse repetition frequencies on forest canopy metrics and biophysical stand properties derived from small-footprint airborne laser data

Source: Elsevier BV

Estimating forest biomass and volume using airborne laser data

Source: Elsevier BV

Estimation of standing wood volume in forest compartments by exploiting airborne laser scanning information: model-based, design-based, and hybrid perspectives

Source: Canadian Science Publishing

Stepwise modelling in ecology and behaviour

Source: Wiley

Accuracy of forest inventory using airborne laser scanning: evaluating the first nordic full-scale operational project

Source: Informa UK Limited

Estim

Source

Fusio

chara

Source

Estim

S

Ac

Source

Speci

using

Source

Deter

Source: Elsevier BV



Regional aboveground forest biomass using airborne and spaceborne LiDAR in Québec.

Source: Elsevier BV

Using airborne laser scanning to monitor tree migration in the boreal–alpine transition zone

Source: Elsevier BV

Mapping stand-level forest biophysical variables for a mixedwood boreal forest using lidar: an examination of scanning density

Source: Canadian Science Publishing

Prediction of tree height, basal area and stem volume in forest stands using airborne laser scanning

Source: Informa UK Limited

Biomass estimation in Neotropical forests using lidar

Source: Wiley

Delineation of Individual Tree Crowns from ALS and Hyperspectral data: a comparison among four methods

Source: Informa UK Limited

Separating the ground and airborne laser sampling phases to estimate tropical forest basal area, volume, and biomass.

Source: Elsevier BV

Quantifying forest above ground carbon content using LiDAR remote sensing

Source: Elsevier BV

Comparing stand inventories for large areas based on photo-interpretation and laser scanning by means of cost-plus-loss analyses

Source: Informa UK Limited

Reducing Uncertainty in Mapping of Mangrove Aboveground Biomass Using Airborne Discrete Return Lidar Data

Source

Predic

two-s

Source

Accur

Source

A

So

Asses

plot s

Source

Effect

growi

Source: Canadian Science Publishing



Practical large-scale forest stand inventory using a small-footprint airborne scanning laser

Source: Informa UK Limited

Influence of Vegetation, Slope, and Lidar Sampling Angle on DEM Accuracy

Source: American Society for Photogrammetry and Remote Sensing

Sensitivity of large-footprint lidar to canopy structure and biomass in a neotropical rainforest

Source: Elsevier BV

Weibull and percentile models for lidar-based estimation of basal area distribution

Source: Informa UK Limited

Area-Based Inventory in Norway – From Innovation to an Operational Reality

Source: Springer Netherlands

Measuring biomass and carbon in delaware using an airborne profiling LIDAR

Source: Informa UK Limited

Error assessment in two lidar-derived TIN datasets

Source: American Society for Photogrammetry and Remote Sensing

Quantifying canopy height underestimation by laser pulse penetration in small-footprint airborne laser scanning data

Source: Informa UK Limited

Functions for Biomass Estimation of Young *Pinus sylvestris*, *Picea abies* and *Betula* spp. from Stands in Northern Sweden with High Stand Densities

Source: Informa UK Limited

Airborne laser scanning—an introduction and overview

Source: Elsevier BV

Effect of flying altitude and pulse repetition frequency on laser scanner penetration rate for digital elevation model generation in a tropical forest

Source

Error

carbo

Source

An Ar

Source

L

So

Meth

Source

Linkin



# Related research

People also read

Recommended articles

Cited by  
59

## Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

## Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

## Open access


- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

## Help and information

- Help and contact
- Newsroom
- All journals
- Books

## Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright © 2024 Taylor & Francis Group, an informa business

Access  
Register  
5 Howick Pl

