

627 62 Views CrossRef citations to date 0 Altmetric

Original Articles

Estimating above-ground biomass in young forests with airborne laser scanning

Erik Næsset 

Pages 473-501 | Received 29 Apr 2009, Accepted 30 Sep 2009, Published online: 06 Feb 2011

 Cite this article  <https://doi.org/10.1080/01431160903474970>

Sample our
Geography
Journals
>> **Sign in here** to start your access
to the latest two volumes for 14 days

 Full Article  Figures & data  References  Citations  Metrics

 Reprints & Permissions  Read this article

We Care About Your Privacy

We and our 855 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting "I Accept" enables tracking technologies to support the purposes shown under "we and our partners process data to provide," whereas selecting "Reject All" or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the ["privacy preferences"] link on the bottom of the webpage [or the floating icon on the bottom-left of the webpage, if applicable]. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. [Here](#)

We and our partners process data to provide:

...

 I Accept

Reject All

Show Purpose



variability of the estimated models ranged from 60% to 83%. Tree species had a significant influence on the models. For given values of the ALS-derived metrics related to canopy height and canopy density, spruce tended to have higher above-ground biomass values than pine and deciduous species. There were no clear effects of model form and canopy threshold on the accuracy of predictions produced by cross validation of the various models, but there is a risk of heteroskedasticity with linear models. Cross validation revealed an accuracy of the root mean square error (RMSE) ranging from 3.85 to 13.9 Mg ha⁻¹, corresponding to 22.6% to 48.1% of mean field-measured biomass. It was concluded that airborne laser scanning has a potential for predicting biomass in young forest stands (> 0.5 ha) with an accuracy of 20–30% of mean ground value.

Related Research Data

Airborne laser scanning as a method in operational forest inventory: Status of accuracy assessments accomplished in Scandinavia

Source: Informa UK Limited

Asses... in the
Field

Source...
Estim... borne
laser

Source...
Estim... direct
latitu



A Mul...
Source...
Biom

Source... rest
Estim...
zone

Source

Estimation of above ground forest biomass from airborne discrete return laser scanner data using canopy-based quantile estimators

Source: Informa UK Limited

A Heteroskedasticity-Consistent Covariance Matrix Estimator and a Direct Test for Heteroskedasticity

Source: JSTOR

Surface lidar remote sensing of basal area and biomass in deciduous forests of eastern Maryland, USA

Source: Elsevier BV

Use of Large-Footprint Scanning Airborne Lidar To Estimate Forest Stand Characteristics in the Western Cascades of Oregon

Source: Elsevier BV

Effects of different sensors, flying altitudes, and pulse repetition frequencies on forest canopy metrics and biophysical stand properties derived from small-footprint airborne laser data

Source: Elsevier BV

Estimating forest biomass and volume using airborne laser data

Source: Elsevier BV

Estimation of standing wood volume in forest compartments by exploiting airborne laser scanning information: model-based, design-based, and hybrid perspectives

Source: Canadian Science Publishing

Stepwise modelling in ecology and behaviour

Source:

Accuracy of forest biomass estimation using airborne laser scanning data in a boreal forest: a full-scale comparison

Source:

Estim

Source:

Fusio

chara

S

Es

Source:

Accu

Source:

Speci

using

Source:

Deter

Source: Elsevier BV



Regional aboveground forest biomass using airborne and spaceborne LiDAR in Québec.

Source: Elsevier BV

Using airborne laser scanning to monitor tree migration in the boreal–alpine transition zone

Source: Elsevier BV

Mapping stand-level forest biophysical variables for a mixedwood boreal forest using lidar: an examination of scanning density

Source: Canadian Science Publishing

Prediction of tree height, basal area and stem volume in forest stands using airborne laser scanning

Source: Informa UK Limited

Biomass estimation in Neotropical forests using lidar

Source: Wiley

Delineation of Individual Tree Crowns from ALS and Hyperspectral data: a comparison among four methods

Source: Informa UK Limited

Separating the ground and airborne laser sampling phases to estimate tropical forest basal area, volume, and biomass.

Source: Elsevier BV

Quantifying forest above ground carbon content using LiDAR remote sensing

Source: Elsevier BV

Comparing stand inventories for large areas based on photo-interpretation and laser scanning

Source:

Reducing uncertainty in forest carbon stocks using airborne laser scanning

Discr

Source:

Predicting forest carbon stocks using airborne laser scanning and optical remote sensing

two-s

Source:

Airborne laser scanning for forest carbon budget estimation

So

Airbo

Source:

Assessing the impact of forest management on carbon stocks using airborne laser scanning

plot scale and landscape scale: a comparison of data sources

Source:

Effect of forest management on carbon stocks: a comparison of forest inventory data and airborne laser scanning data

Source: Canadian Science Publishing

Practical large-scale forest stand inventory using a small-footprint airborne scanning laser

Source: Informa UK Limited

Influence of Vegetation, Slope, and Lidar Sampling Angle on DEM Accuracy

Source: American Society for Photogrammetry and Remote Sensing

Sensitivity of large-footprint lidar to canopy structure and biomass in a neotropical rainforest

Source: Elsevier BV

Weibull and percentile models for lidar-based estimation of basal area distribution

Source: Informa UK Limited

Area-Based Inventory in Norway – From Innovation to an Operational Reality

Source: Springer Netherlands

Measuring biomass and carbon in delaware using an airborne profiling LIDAR

Source: Informa UK Limited

Error assessment in two lidar-derived TIN datasets

Source: American Society for Photogrammetry and Remote Sensing

Quantifying canopy height underestimation by laser pulse penetration in small-footprint airborne laser scanning data

Source: Informa UK Limited

Functions for Biomass Estimation of Young *Pinus sylvestris*, *Picea abies* and *Betula* spp. from Stands in Northern Sweden with High Stand Densities

Source: Informa UK Limited

Airbo

Source

Effect

rate f

Source

Error

carbo

Source

A

So

LiDAR

Source

Meth

Source

Linkin



Related research 

- People also read
- Recommended articles
- Cited by 62

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access



- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up

Register to receive updates by email



Elsevier
an Elsevier company