



678 | 67

Views | CrossRef citations to date | 0 Altmetric

Original Articles

Estimating above-ground biomass in young forests with airborne laser scanning

Erik Næsset

Pages 473-501 | Received 29 Apr 2009, Accepted 30 Sep 2009, Published online: 06 Feb 2011

Cite this article <https://doi.org/10.1080/01431160903474970>

Sample our
Geography
Journals
>> **Sign in here** to start your access
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

Read this article

Share

Abstract

Total above-ground biomass of spruce, pine and birch was estimated in three different field datasets collected in young forests in south-east Norway. The mean heights ranged from 1.77 to 9.66 m. These field data were regressed against metrics derived from canopy height distributions generated from airborne laser scanner (ALS) data with a point density of 0.9–1.2 m⁻². The field data consisted of 79 plots with size 200–232.9 m² and 20 stands with an average size of 3742 m². Total above-ground biomass ranged from 2.27 to 90.42 Mg ha⁻¹. The influences of (1) regression model form, (2) canopy threshold value and (3) tree species on the relationships between biomass and ALS-derived metrics were assessed. The analysed model forms were multiple linear models, models with logarithmic transformation of the response and explanatory variables, and models with square root transformation of the response. The different canopy

thresholds considered were fixed values of 0.5, 1.3 and 2.0 m defining the limit between laser canopy echoes and below-canopy echoes. The proportion of explained variability of the estimated models ranged from 60% to 83%. Tree species had a significant influence on the models. For given values of the ALS-derived metrics related to canopy height and canopy density, spruce tended to have higher above-ground biomass values than pine and deciduous species. There were no clear effects of model form and canopy threshold on the accuracy of predictions produced by cross validation of the various models, but there is a risk of heteroskedasticity with linear models. Cross validation revealed an accuracy of the root mean square error (RMSE) ranging from 3.85 to 13.9 Mg ha⁻¹, corresponding to 22.6% to 48.1% of mean field-measured biomass. It was concluded that airborne laser scanning has a potential for predicting biomass in young forest stands (> 0.5 ha) with an accuracy of 20–30% of mean ground value.

Related Research Data

[Airborne laser scanning as a method in operational forest inventory: Status of accuracy assessments accomplished in Scandinavia](#)

Source: Scandinavian Journal of Forest Research

[Assessing Point Accuracy of DGPS Under Forest Canopy Before Data Acquisition, in the Field and After Postprocessing](#)

Source: Scandinavian Journal of Forest Research

[Estimating tree heights and number of stems in young forest stands using airborne laser scanner data](#)

Source: Remote Sensing of Environment

[A Multiple Resource Inventory of Delaware Using Airborne Laser Data](#)

Source: BioScience

[Lidar remote sensing of above-ground biomass in three biomes](#)

Source: Global Ecology and Biogeography

[Estimation of above- and below-ground biomass across regions of the boreal forest zone using airborne laser](#)

People also read

Recommended articles

Cited by
67

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up

