

484 Views | 18 CrossRef citations to date | 1 Altmetric

Miscellany

The expected child versus the actual child: implications for the mother–baby bond

H. Pearce & S. Ayers 

Pages 89-102 | Received 24 Mar 2002, Accepted 20 Feb 2004, Published online: 23 Jan 2007

🗨️ Cite this article [🔗 https://doi.org/10.1080/0264683042000325528](https://doi.org/10.1080/0264683042000325528)

Sample our Behavioral Sciences Journals 
>> **Sign in here** to start your access to the latest two volumes for 14 days

 Full Article  Figures & data  References  Citations  Metrics

 Reprints & Permissions

[Read this article](#)

[Share](#)

Abstract

The aim of this study was to examine whether women's perceptions of their infant remain stable between late pregnancy and the early postnatal period, and to examine the effect of women's expectations and evaluations of their infant on the mother–baby bond. Participants completed questionnaires at 39 weeks gestation and 3 weeks after birth. Questionnaires measured demographic details, expected/actual infant characteristics, symptoms of anxiety and depression and the mother–baby bond (postpartum only). It was found that there was a significant difference between expectations and evaluations, with infants being perceived as less fussy, more adaptable and dull than they were expected to be. Both expectations of the baby and evaluation of the baby had a significant effect on the mother–baby bond. Women who expected their infant to have a more difficult temperament and then evaluated them,

similarly reported a poorer mother–baby bond postpartum. Symptoms of anxiety and depression in pregnancy were unrelated to expectations of the baby, or to the postnatal mother–baby bond. The effects of parity and postnatal anxiety and depression were also examined. The results are discussed, suggesting that a subgroup of women who have negative perceptions of their infant before and after birth may be at risk of mother–infant attachment problems and long-term mother–infant difficulties.

Notes

ns = nonsignificant.

Possible range: Fussy 9–63; Unadaptable 5–35; Dull 4–28; Unpredictable 5–35; Total 23–161.

Related Research Data

[A validation study of the Hospital Anxiety and Depression Scale \(HADS\) in different groups of Dutch subjects](#)

Source: Psychological Medicine

[Social support, self-esteem and depression](#)

Source: Psychological Medicine

[Newborn Behavior and Risk of Postnatal Depression in the Mother](#)

Source: Infancy

[Social Support, Infant Temperament, and Parenting Self-Efficacy: A Mediational Model of Postpartum Depression](#)

Source: Child Development

[Anxiety and the Allocation of Attention to Threat](#)

Source: The Quarterly Journal of Experimental Psychology Section A

[The Hospital Anxiety and Depression Scale](#)

Source: Acta Psychiatrica Scandinavica

[Measurement of Infant Difficultiness](#)



Related research

People also read

Recommended articles

Cited by
18

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG

 Taylor and Francis
Group