Taylor & Francis
Online

Food Additives & Contaminants ›
Volume 21, 2004 - Issue 5

161 Views | 43 CrossRef citations to date | 0 Altmetric

Original Articles

Occurrence of aflatoxin M_1 in domestic milk in Japan during the winter season

M. Nakajima, S. Tabata, H. Akiyama, Y. Itoh, T. Tanaka, H. Sunagawa, ...show all

Pages 472-478 | Received 05 Sep 2003, Accepted 18 Feb 2004, Published online: 20 Feb 2007

Cite this article https://doi.org/10.1080/02652030410001677817

Sample our
Food Science & Technology
Journals
>> Sign in here to start your access
to the latest two volumes for 14 days

Full Article Figures & data References Citations Metrics
Reprints & Permissions Read this article Share

Abstract

A total of 208 samples of commercial pasteurized milk gathered from retail outlets across Japan during the winter season were analysed for aflatoxin M_1 (AFM_1). Japan was divided into 11 regions from north to south, and nine to 45 milk samples from each region were randomly purchased between December 2001 and February 2002. Each milk sample was cleaned up by an immunoaffinity column, and AFM_1 was quantified by liquid chromatography with fluorescence detection in four independent laboratories. The limit of detection of the method was 0.001 μg kg^{-1}. The identity of the putative AFM_1 in milk sample was confirmed by the formation of AFM_1 hemi-acetal with trifluoroacetic acid. Based on the results obtained with spiked samples (0.05 μg AFM_1 kg^{-1}), the mean recovery was 91.4%, the relative standard deviation for repeatability was 4.6%, and the relative standard deviation for reproducibility was 8.0% among four

independent laboratories. AFM_1 was detected in 207 (99.5%) of 208 milk samples at 0.001–0.029 μg kg^{-1}, with a mean of 0.009 μg kg^{-1} and a 90th percentile of 0.014 μg kg^{-1}. No significant difference of the level of AFM_1 contamination was observed among the regions.

Keywords:

survey | aflatoxin M1 | milk | immunoaffinity column | liquid chromatography

Acknowledgement

Work was supported in part by a grant for Scientific Research Expense for Health and Welfare Programs from the Japanese Government.

Related Research Data

Investigation of aflatoxins contamination in foods and foodstuffs.
Source: Food Hygiene and Safety Science (Shokuhin Eiseigaku Zasshi)
Reverse Phase Liquid Chromatographic Determination and Confirmation of Aflatoxin M1 in Cheese
Source: Journal of AOAC INTERNATIONAL
Natural occurrence of aflatoxin M1 in imported and domestic cheese. Studies on mycotoxins in foods. VI.
Source: Food Hygiene and Safety Science (Shokuhin Eiseigaku Zasshi)
Cyclopiazonic acid in combination with aflatoxins, zearalenone and ochratoxin A in Indonesian corn
Source: Mycopathologia
Occurrence of mycotoxins in raw ingredients used for animal feeding stuffs in the United Kingdom in 1992
Source: Food Additives & Contaminants
Determination of aflatoxin M1in milk by reversed-phase high-performance liquid

Related research

People also read | Recommended articles | Cited by 43

Exposure to aflatoxins in Japan: risk assessment for aflatoxin B1 >

Y. Sugita-Konishi et al.
Food Additives & Contaminants: Part A
Published online: 5 Jan 2010

Information for

Authors
R&D professionals
Editors
Librarians
Societies

Open access

Overview
Open journals
Open Select
Dove Medical Press
F1000Research

Opportunities

Reprints and e-prints
Advertising solutions
Accelerated publication
Corporate access solutions

Help and information

Help and contact
Newsroom
All journals
Books

Keep up to date

Register to receive personalised research and resources by email

Sign me up

Copyright © 2026 Informa UK Limited Privacy policy
Cookies Terms & conditions Accessibility

Taylor & Francis by informa

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG