


Food Additives & Contaminants >

Volume 23, 2006 - [Issue 2](#)

341 | 70 | 0  
Views | CrossRef citations to date | Altmetric

Original Articles

# Distribution and stability of Aflatoxin M<sub>1</sub> during processing and ripening of traditional white pickled cheese

H. H. Oruc , R. Cibik, E. Yilmaz & O. Kalkanli

Pages 190-195 | Received 10 Aug 2005, Accepted 20 Oct 2005, Published online: 11 Feb 2011

 Cite this article  <https://doi.org/10.1080/02652030500389048>

Sample our  
Environment & Agriculture  
Journals  
>> [Sign in here](#) to start your access  
to the latest two volumes for 14 days

 Full Article

 Figures & data

 References

 Citations

 Metrics

 Reprints & Permissions

Read this article

Share

## Abstract

The distribution of aflatoxin M<sub>1</sub> (AFM<sub>1</sub>) has been studied between curd, whey, cheese and pickle samples of Turkish white pickled cheese produced according to traditional techniques and its stability studied during the ripening period. Cheeses were produced in three cheese-making trials using raw milk that was artificially contaminated with AFM<sub>1</sub> at the levels of 50, 250 and 750 ng/l and allowed to ripen for three months. AFM<sub>1</sub> determinations were carried out at intervals by LC with fluorescence detection after immunoaffinity column clean-up. During the syneresis of the cheese a proportionately high concentration of AFM<sub>1</sub> remained in curd and for each trial the level was 3.6, 3.8 and 4.0 times higher than levels in milk. At the end of the ripening, the distribution of AFM<sub>1</sub> for cheese/whey + brine samples was 0.9, 1.0 and 1.3 for first, second and third

spiking respectively indicating that nearly half of the AFM<sub>1</sub> remained in cheese. It has been found that only 2–4% of the initial spiking of AFM<sub>1</sub> transferred into the brine solution. During the ripening period AFM<sub>1</sub> levels remained constant suggesting that AFM<sub>1</sub> was quite stable during manufacturing and ripening.

Keywords:

Aflatoxin M<sub>1</sub> traditional white pickled cheese stability distribution

## Acknowledgments

This research was funded by the Uludag University Research Funds (project no 2002/73).

### Related Research Data

[Fate of Aflatoxin M1 in Brick and Limburger-like Cheese](#)

Source: Journal of Food Protection

[Fate of Aflatoxin M1 in Cheddar Cheese and in Process Cheese Spread](#)

Source: Journal of Food Protection

[Occurrence and Stability of Aflatoxin M1 in Milk and Milk Products: A Worldwide Review](#)

Source: Journal of Food Protection

[Verhalten von Aflatoxin M1 während der Reifung und Lagerung von Käse](#)

Source: European Food Research and Technology

[Ability of Dairy Strains of Lactic Acid Bacteria to Bind Aflatoxin M1 in a Food Model](#)

Source: Journal of Food Protection

[Fate of Aflatoxin M1 in Parmesan and Mozzarella Cheese](#)

Source: Journal of Food Protection

[Detection of aflatoxin M1 in cheese samples by ELISA](#)

Source: Food Control

[Principles of Analytical Chemistry](#)

People also read

Recommended articles

Cited by  
70

From Cow's Milk to Cheese, Yogurt, and Labneh: Evaluating aflatoxin M1 fate in Traditional Lebanese Dairy Processing and the Efficacy of Regulations through a R... >

Rouaa Daou et al.

Food Additives & Contaminants: Part A

Published online: 11 Mar 2025

## Information for

Authors

R&D professionals

Editors

Librarians

Societies

## Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

## Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

## Help and information

Help and contact

Newsroom

All journals

Books

## Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954  
5 Howick Place | London | SW1P 1WG



Taylor & Francis  
by informa