Home ▶ All Journals ▶ Journal of Applied Statistics ▶ List of Issues ▶ Volume 43, Issue 10 Collinearity diagnostic applied in ridge

Journal of Applied Statistics > Volume 43, 2016 - Issue 10

1.845 97

Views CrossRef citations to date Altmetric

Articles

Collinearity diagnostic applied in ridge estimation through the variance inflation factor

Roman Salmerón Gómez, José García Pérez, María Del Mar López Martín & Catalina García García 🔀

Pages 1831-1849 | Received 22 Apr 2015, Accepted 12 Nov 2015, Published online: 25 Feb 2016

66 Cite this article

▶ https://doi.org/10.1080/02664763.2015.1120712





Full Article

Figures & data

References

66 Citations

Metrics

Reprints & Permissions

Read this article

ABSTRACT

The variance inflation factor (VIF) is used to detect the presence of linear relationships between two or more independent variables (i.e. collinearity) in the multiple linear regression model. However, the traditionally used VIF definitions encounter some

problem an exter overcom express

Q KEYWO

About Cookies On This Site

We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our Privacy Policy

er presents Accept All Essential Onlyn, The endent Settings

Acknowledgments

We would like to thank referees for their detailed, encouraging and constructive reviews of our paper which clearly contributed to improving both the structure and the content of this manuscript.

Disclosure statement


No potential conflict of interest was reported by the authors.

ORCID

Catalina García García http://orcid.org/0000-0003-1622-3877

Notes

1. Note that a standardized variable is the value of the variable minus its mean, divided by the square root of the number of observations multiplied by its variance while a typified variable is the value of the variable minus its mean, divided by its standard deviation.



Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up











Copyright © 2024 Informa UK Limited Privacy policy Cookies Terms & conditions



Accessibility

Registered in England & Wales No. 3099067 5 Howick Place | London | SW1P 1WG

About Cookies On This Site



We and our partners use cookies to enhance your website experience, learn how our site is used, offer personalised features, measure the effectiveness of our services, and tailor content and ads to your interests while you navigate on the web or interact with us across devices. You can choose to accept all of these cookies or only essential cookies. To learn more or manage your preferences, click "Settings". For further information about the data we collect from you, please see our Privacy Policy



Essential Onl

Settings