



Journal of Applied Statistics >

Volume 43, 2016 - [Issue 10](#)

2,186	133	0
Views	CrossRef citations to date	Altmetric

Articles

# Collinearity diagnostic applied in ridge estimation through the variance inflation factor

Roman Salmerón Gómez, José García Pérez, María Del Mar López Martín & Catalina García García

Pages 1831-1849 | Received 22 Apr 2015, Accepted 12 Nov 2015, Published online: 25 Feb 2016

Cite this article <https://doi.org/10.1080/02664763.2015.1120712>



Sample our  
Mathematics & Statistics  
Journals



>> **Sign in here** to start your access  
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

Read this article

Share

## ABSTRACT

The variance inflation factor (VIF) is used to detect the presence of linear relationships between two or more independent variables (i.e. collinearity) in the multiple linear regression model. However, the traditionally used VIF definitions encounter some problems when extended to the case of the ridge estimation (RE). This paper presents an extension of the VIF in RE by providing two alternative VIF expressions that overcome these problems in the general case. Some characteristics of these expressions are also presented and compared with the traditional expression. The results are illustrated with an economic example in the case of three independent variables and with a Monte Carlo simulation for the general case.

KEYWORDS:

## Acknowledgments

We would like to thank referees for their detailed, encouraging and constructive reviews of our paper which clearly contributed to improving both the structure and the content of this manuscript.

---

## Disclosure statement

No potential conflict of interest was reported by the authors.

---

## ORCID

Catalina García García <http://orcid.org/0000-0003-1622-3877>

---

## Notes

1. Note that a standardized variable is the value of the variable minus its mean, divided by the square root of the number of observations multiplied by its variance while a typified variable is the value of the variable minus its mean, divided by its standard deviation.

---

## Related research

People also read

Recommended articles

Cited by  
133

## Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

## Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

## Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

## Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

## Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)

 Taylor and Francis Group

Registered in England & Wales No. 01072954  
5 Howick Place | London | SW1P 1WG