

Brain Injury >

Volume 22, 2008 - [Issue 6](#)

583 Views | 57 CrossRef citations to date | 50 Altmetric

Original

# Ten-year outcomes following traumatic brain injury: A population-based cohort

C. M. Cameron, D. M. Purdie, E. V. Kliewer & Professor R. J. McClure Director 

Pages 437-449 | Received 12 Nov 2007, Accepted 17 Mar 2008, Published online: 03 Jul 2009

 Cite this article  <https://doi.org/10.1080/02699050802060621>

Sample our  
Medicine, Dentistry, Nursing  
& Allied Health Journals  
>> **Sign in here** to start your access  
to the latest two volumes for 14 days

 Full Article  Figures & data  References  Citations  Metrics

 Reprints & Permissions

[Read this article](#)

 Share

## Abstract

**Primary objective:** To quantify the 10 year health service use (HSU) and mortality outcomes for people with a traumatic brain injury (TBI).

**Research design:** A population-based matched cohort study using linked administrative data from Manitoba, Canada (Manitoba Injury Outcome Study).

**Methods and procedures:** An inception cohort (1988–1991) of hospitalized cases with TBI aged 18–64 years (n= 1290) was identified and matched to a non-injured comparison group (n= 1290). Survival analysis, Negative binomial and Poisson regression were used to quantify associations between injury and HSU/mortality outcomes for 10 years following the TBI event.

Main outcome and results: The majority of deaths (47.2%) occurred in the first 60 days following injury. Excluding the first 60 days, the adjusted 10 year mortality remained elevated (mortality rate ratio = 1.48, 95% CI = 1.02-2.15). After adjusting for demographic characteristics and pre-existing health status, the TBI cohort had more post-injury hospitalizations (rate ratio (RR) = 1.54, 95% CI = 1.39-1.71), greater cumulative lengths of stay (RR = 5.14, 95% CI = 3.29-8.02) and a greater post-injury physician claims rate (RR = 1.44, 95% CI = 1.35-1.53) than the non-injured cohort.

Conclusions: People who sustain a TBI and survive the initial acute phase of care experience substantially increased long-term morbidity compared to the general population, regardless of the level of injury severity.

Epidemiology

traumatic brain injury

data linkage

outcomes

mortality

morbidity

health service utilization

## Related Research Data

### [Measuring Disability and Quality of Life Postinjury](#)

Source: Unknown Repository

### [Functional outcome after rehabilitation for severe traumatic brain injury](#)

Source: Archives of Physical Medicine and Rehabilitation

### [Return to productivity following traumatic brain injury: Cognitive, psychological, physical, spiritual, and environmental correlates](#)

Source: Disability and Rehabilitation

### [Understanding outcomes based on the post-acute hospitalization pathways followed by persons with traumatic brain injury](#)

Source: Brain Injury

### [Injury Hospitalization and Risks for Subsequent Self-Injury and Suicide: A National Study From New Zealand](#)

Source: American Journal of Public Health

### [Secondary conditions following spinal cord injury in a population-based sample](#)

Source: Spinal Cord

## Related research

People also read

Recommended articles

Cited by  
57

### Information for

Authors  
R&D professionals  
Editors  
Librarians  
Societies

### Opportunities

Reprints and e-prints  
Advertising solutions  
Accelerated publication  
Corporate access solutions

### Open access

Overview  
Open journals  
Open Select  
Dove Medical Press  
F1000Research

### Help and information

Help and contact  
Newsroom  
All journals  
Books

### Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954  
5 Howick Place | London | SW1P 1WG

 **Taylor & Francis**  
by informa