

Brain Injury >

Volume 22, 2008 - [Issue 6](#)

575 Views | 55 CrossRef citations to date | 50 Altmetric

Original

Ten-year outcomes following traumatic brain injury: A population-based cohort

C. M. Cameron, D. M. Purdie, E. V. Kliewer & Professor R. J. McClure Director 

Pages 437-449 | Received 12 Nov 2007, Accepted 17 Mar 2008, Published online: 03 Jul 2009

 Cite this article  <https://doi.org/10.1080/02699050802060621>

Sample our
Medicine, Dentistry, Nursing
& Allied Health Journals
>> **Sign in here** to start your access
to the latest two volumes for 14 days



 Full Article

 Figures & data

 References

 Citations

 Metrics

 Reprints & Permissions

[Read this article](#)

 Share

Abstract

Primary objective: To quantify the 10 year health service use (HSU) and mortality outcomes for people with a traumatic brain injury (TBI).

Research design: A population-based matched cohort study using linked administrative data from Manitoba, Canada (Manitoba Injury Outcome Study).

Methods and procedures: An inception cohort (1988–1991) of hospitalized cases with TBI aged 18–64 years (n= 1290) was identified and matched to a non-injured comparison group (n= 1290). Survival analysis, Negative binomial and Poisson regression were used to quantify associations between injury and HSU/mortality outcomes for 10 years following the TBI event.

Main outcome and results: The majority of deaths (47.2%) occurred in the first 60 days following injury. Excluding the first 60 days, the adjusted 10 year mortality remained elevated (mortality rate ratio = 1.48, 95% CI = 1.02-2.15). After adjusting for demographic characteristics and pre-existing health status, the TBI cohort had more post-injury hospitalizations (rate ratio (RR) = 1.54, 95% CI = 1.39-1.71), greater cumulative lengths of stay (RR = 5.14, 95% CI = 3.29-8.02) and a greater post-injury physician claims rate (RR = 1.44, 95% CI = 1.35-1.53) than the non-injured cohort.

Conclusions: People who sustain a TBI and survive the initial acute phase of care experience substantially increased long-term morbidity compared to the general population, regardless of the level of injury severity.

Epidemiology

traumatic brain injury

data linkage

outcomes

mortality

morbidity

health service utilization

Related Research Data

[Measuring Disability and Quality of Life Postinjury](#)

Source: Unknown Repository

[Functional outcome after rehabilitation for severe traumatic brain injury](#)

Source: Archives of Physical Medicine and Rehabilitation

[Return to productivity following traumatic brain injury: Cognitive, psychological, physical, spiritual, and environmental correlates](#)

Source: Disability and Rehabilitation

[Understanding outcomes based on the post-acute hospitalization pathways followed by persons with traumatic brain injury](#)

Source: Brain Injury

[Injury Hospitalization and Risks for Subsequent Self-Injury and Suicide: A National Study From New Zealand](#)

Source: American Journal of Public Health

[Secondary conditions following spinal cord injury in a population-based sample](#)

Source: Spinal Cord

Related research

People also read

Recommended articles

Cited by
55

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)

 Taylor and Francis Group

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG