

241 | 5 | 0
Views | CrossRef citations to date | Altmetric

Original Articles

Analytical study on ethephon residue determination in water by ion-pairing liquid chromatography/tandem mass spectrometry

Cristina Ripollés, José M. Marín, Juan V. Sancho, Francisco J. López & Félix Hernández

Pages 1380-1391 | Received 25 May 2010, Accepted 27 Aug 2010, Published online: 21 Oct 2011

🗨️ Cite this article 🔗 <https://doi.org/10.1080/03067319.2010.520131>

Sample our
Environment and Sustainability
Journals

>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

📄 Full Article

🖼️ Figures & data

📖 References

🗨️ Citations

📊 Metrics

🖨️ Reprints & Permissions

Read this article

🔗 Share

Abstract

A detailed analytical study on ethephon residue determination in water, making use of ion-pairing liquid chromatography coupled to electrospray tandem mass spectrometry (LC/MS/MS), has been carried out. Ethephon is a plant growth regulator, highly polar, which is typically present in aqueous solution in anionic form due to its acid character. Both its extraction and pre-concentration from water samples and its chromatographic retention are difficult. Several approaches for sample pretreatment have been tested including direct injection into the chromatographic system, on-line solid phase extraction (SPE) and off-line SPE, with the best results being obtained after off-line SPE, using Oasis MAX cartridges (mixed-mode strong anion-exchange). After testing several ion-pairing reagents, tetrabutylammonium acetate (TBA) was selected. This was

added to the samples before LC/MS/MS analysis to facilitate ethephon chromatographic retention. The acquisition of several specific MS/MS transitions together with the evaluation of their relative intensity ratios allowed the reliable confirmation of the analyte in samples. The optimised approach was tested in low-salinity water spiked at $0.1 \mu\text{g L}^{-1}$ level with satisfactory recovery, and a limit of detection of $0.02 \mu\text{g L}^{-1}$. To this purpose, the water sample was partially de-ionised in an initial stage, in order to remove major ions that would have interfered in analyses. The application of this methodology to more saline/complex water samples, as surface or wastewater, was problematic and a thorough optimisation of the de-ionisation conditions would be required.

Keywords:

ethephon ion-pairing liquid chromatography tandem mass spectrometry
tetrabutylammonium, water analysis

Acknowledgements

This work has been developed under financial support of the Ministry of Education and Science, Spain (CTM2006-06417). The authors acknowledge the financial support of Generalitat Valenciana, as research group of excellence PROMETEO/2009/054. C. Ripollés is very grateful to the Ministry of Education and Science for her pre-doctoral grant.

Related research

People also read

Recommended articles

Cited by
5

[Determination of Ethephon in Pesticide Formulations by Ion Exchange Chromatography with Indirect Spectrophotometric Detection >](#)

Niki C. Maragou et al.
Analytical Letters
Published online: 21 Oct 2019

Information for

[Authors](#)
[R&D professionals](#)
[Editors](#)
[Librarians](#)
[Societies](#)

Opportunities

[Reprints and e-prints](#)
[Advertising solutions](#)
[Accelerated publication](#)
[Corporate access solutions](#)

Keep up to date

Register to receive personalised research and resources by email

 [Sign me up](#)



Open access

[Overview](#)
[Open journals](#)
[Open Select](#)
[Dove Medical Press](#)
[F1000Research](#)

Help and information

[Help and contact](#)
[Newsroom](#)
[All journals](#)
[Books](#)

Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG



Taylor & Francis
by informa