

Free access

1,623 Views | 12 CrossRef citations to date | 0 Altmetric

Listen

Original Articles

Nutritional secondary hyperparathyroidism and osteodystrophia fibrosa in a Hodgson's hawk-eagle (*Spizaetus nipalensis*)

T. Toyoda, K. Ochiai, M. Komatsu, T. Kimura & T. Umemura

Pages 9-12 | Published online: 08 Jun 2010

Cite this article <https://doi.org/10.1080/03079450310001636237>

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

View PDF

Abstract

A Hodgson's hawk-eagle (*Spizaetus nipalensis*) reared by a falconer showed severe weakness and emaciation. The bird was fed a diet consisting of 100% meat. Serological examination of the blood revealed hypercalcaemia, hyperphosphataemia, severe osteopenia, and bone resorption. The bone resorption contained hypercalcaemia, hyperphosphataemia, and hyperparathyroidism.

We Care About Your Privacy

We and our 842 partners store and/or access information on a device, such as unique IDs in cookies to process personal data. You may accept or manage your choices by clicking below, including your right to object where legitimate interest is used, or at any time in the privacy policy page. These choices will be signaled to our partners and will not affect browsing data. [Privacy Policy](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device characteristics for identification. Store and/or access information on a device. Personalised advertising and content, advertising and content measurement, audience research and services development.

List of Partners (vendors)

I Accept

Essential Only

Show Purpose

Nutritional secondary hyperparathyroidism caused by long term deficiency of vitamin D₃, calcium or phosphorus, or by imbalances in the last two nutrients causes excessive production of parathyroid hormone (PTH), resulting in osteoclastic bone absorption (Woodard, [1997](#)). Rickets, which has the same pathogenesis, occurs in various kinds of young birds and mammals and is well described (Whitehead & Wilson, [1992](#); Thorp, [1994](#)). However, description of osteodystrophia fibrosa (fibrous osteodystrophy) in avian species is limited (Long et al., [1983](#)). Here, we describe the disease secondary to nutritional hyperparathyroidism in a Hodgson's hawk-eagle (*Spizaetus nipalensis*) bred for falconry.

2 Materials and Methods

Necropsy was performed and the liver, kidneys, heart, lungs, intestine, pancreas, adrenal glands, thyroid glands, parathyroid glands, sternum, femur, tibiotarsus and skeletal muscles were routinely fixed in 10% neutral buffered formalin, embedded in paraffin wax and stained with haematoxylin and eosin. Selected samples of the sternum, femur and tibiotarsus were decalcified by formic acid before embedding.

3 Results

A 12-year-old male Hodgson's hawk-eagle (*Spizaetus nipalensis*) reared by a falconer in Japan was presented with weakness, weight loss and a swollen right leg para-

viscera were examined. The traditional bone fracture of the left tibiotarsus was observed. The long bones of the left tibiotarsus were decalcified and the diaphysis of the long bones was examined. The diaphysis of the left tibiotarsus was apparently




and calcium compound by oral administration. Biochemical examination eight days later revealed lower serum calcium (72.0 µg/ml) and phosphorus (29.0 µg/ml) and higher alkaline phosphatase (ALP) (3100 IU/l) than normal. Normal ranges of these inorganic substances or enzyme in raptors are 89.3–101.9 µg/ml (calcium), 30.3–43.4 µg/ml (phosphorus) and 31–257 IU/l (ALP), respectively (Samour, [2000](#)). The bird died ten days after the initial examination.


Gross examination confirmed that bone was very fragile with multiple complete and/or incomplete fractures. The parathyroid glands were bilaterally enlarged, up to 6 mm in diameter, and they were even larger than thyroid glands.

Histological changes of the affected bone mainly consisted of osteomalacic lesions and fractures. The former was characterized by a significant decrease in the amounts of mineralized trabecular bone associated with an increase in the width of unmineralized osteoid ([Fig. 1](#)). The osteoid borders were occasionally lined by osteoblasts and distinct osteoid seams were recognized. These changes occurred in the cortical and cancellous bone of the sternum, femur and tibiotarsus. The cortical bones of the diaphyses were also markedly thinned with dilation of the Haversian canals. The femur and tibiotarsus partly showed subperiosteal proliferation of trabeculae and loose fibrous tissue ([Fig. 2](#)). In addition, numerous osteoclasts were frequently observed on the surface of the trabeculae, Haversian canals and endosteum and fibrous tissue proliferated between these trabeculae ([Fig. 3](#)), suggesting a progression to osteodystrophia fibrosa. Although there were endochondral and membranous ossifications in and around the necrotic bone of each fracture, mineralization was incomplete in these foci. Enlarged

parathyroid glands had
lightly enlarged
atrophic


Fig. 1 Decreased amounts of unmineralized osteoid
Haversian canals





Display full size

Fig. 2 Decalcified section of the tibiotarsus. Marked periosteal bone reaction with proliferation of osteoid trabeculae and loose fibrous tissue at the exterior of thinning cortex bone (arrowheads). HE. Bar=500 μ m.




Display




Fig. 3 De
of bone
(arrowhe

the surface
cavity



Display full size

Fig. 4 Parathyroid gland showing proliferation of chief cells with cytoplasmic swelling and vacuolation. HE. Bar=50 μ m.



Display full size



Other
moderat
fibrosis.

nal cortex,
cardial

Osteodystrophia fibrosa is the result of continuous and excessive secretion of PTH and is characterized by marked osteoclastic resorption and fibrous replacement (Palmer, [1991](#)). Locomotory disorders with fractures in birds of prey including those for falconry, have been reported as "cramps" or "fits"; protracted dietary deficiency of calcium or imbalance of calcium/phosphorus ratio has been considered important causative factors (Wallach & Flieg, [1970](#); Cooper, [1975](#)). However, pathological descriptions of osteodystrophia fibrosa due to nutritional secondary hyperparathyroidism have rarely been reported in avian species (Long et al., [1983](#)).

The present case was considered nutritional secondary hyperparathyroidism that resulted in severe osteodystrophia fibrosa. The active osteoclastic bone resorption and fibrous tissue proliferation between trabeculae in the examined bones are characteristic findings of osteodystrophia fibrosa. On the other hand, accumulation of osteoid and formation of osteoid seams in the medullary cavity are specific features of osteomalacia, which frequently progresses to osteodystrophia fibrosa when severe hyperparathyroidism develops in the course of the disease (Woodard, [1997](#)). Compensating subperiosteal bone proliferation occurs in some cases of osteopenia (Riddell, [1996](#)). However, subperiosteal proliferation of osteoid and fibrous tissues in the present case was considered as unmineralized reactive callus resulting from microscopic fractures, because it was observed only in part of the femur and tibiotarsus.

The calcium: phosphorus ratio (Ca: P) of meat and fish commonly fed to birds of prey range from 1: 17 to 1: 44 whereas the correct Ca: P ratio for avian diets is listed as 1.5:

1 (Wallach & Flieg, 1970). In addition to an abnormal Ca: P ratio, hyperparathyroidism is characterized by osteoclastic resorption of bone, resulting in osteopenia. Although the type of hyperparathyroidism is important for the



For a long time, falconry has been maintained throughout the world as a traditional form of hunting and the avian species used include hawks, eagles, buzzards and peregrines. Dietary requirements of raptors have not been sufficiently studied, but artificial diets such as muscle meat and eviscerated prey are known to lack vitamins, calcium and phosphorus and may cause various metabolic bone diseases in raptors (Keymer, [1972](#); De Water, [1996](#)). Investigation of the dietary requirements of raptors is essential to preserve these rare birds.

Acknowledgments

The authors would like to thank Dr C. Itakura of the Institute of Physical and Chemical Research, Wako, Japan, for his suggestions.

References

1. Cooper , J.E. 1975 . Osteodystrophy in birds of prey . *Veterinary Record* , 97 : 307

[PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

2. De Water D.V. Raptor rehabilitation, *Diseases of Cage and Aviary Birds* 3rd edn, Roskopf W., Woerpel R. (eds) Williams & Wilkins: Baltimore 1996 1007 1028

[Goog](#)



3. Keymer [1972](#) . *Veterinary Record* , 90 : 594 .

4. Lor [1972](#) . *Veterinary Record* , 90 : 839 -

5. Palme [1972](#) . *Veterinary Record* , 90 : 594 .

Press.

[Google Scholar](#)

6. Riddell C. Skeletal system, Avian Histopathology 2nd edn, Riddell C. (ed.) American Association of Avian Pathologists: Kennet Square, PA 1996 45 60

[Google Scholar](#)

7. Roth , S.I. and Capen , C.C. 1974 . Ultrastructural and functional correlations of the parathyroid gland . International Review of Experimental Pathology , 13 : 161 - 221 .

[PubMed](#) | [Google Scholar](#)

8. Samour, J. (2000). Avian Medicine. London: Harcourt Publishers.

[Google Scholar](#)

9. Thorp , B.H. 1994 . Skeletal disorders in the fowl: a review . Avian Pathology , 23 : 203 - 236 .

 | [PubMed](#) | [Web of Science ®](#) | [Google Scholar](#)

10. Wallach, J.D. & Flieg, G.M. (1970). Cramps and fits in carnivorous birds. In J. Lucas (Ed.), International Zoo Yearbook Volume 10 (pp. 3-4). London: Zoological Society of London.

[Google Scholar](#)

11. White
Skelet
265 2

[Goog](#)

12. Wor
Kir
unt R.D.,

[Goog](#)

Download

Related research

People also read

Recommended articles

Cited by
12

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright © 2024 Taylor & Francis Group, an informa business



Register to receive personalised research and resources by email

5 Howick Place