

23,136 240

Views

CrossRef citations to date

3

Altmetric

Sampling Theory

Estimation Under Purposive Sampling

Jacqueline M. Guarte & Erniel B. Barrios 

Pages 277-284 | Received 01 Sep 2004, Accepted 01 Aug 2005, Published online: 15 Feb 2007

 Cite this article  <https://doi.org/10.1080/03610910600591610>

Sample our
Mathematics & Statistics
Journals

>> [Sign in here](#) to start your access
to the latest two volumes for 14 days



 Full Article

 Figures & data

 References

 Citations

 Metrics

 Reprints & Permissions

Read this article

Share

Abstract

Purposive sampling is described as a random selection of sampling units within the segment of the population with the most information on the characteristic of interest. Nonparametric bootstrap is proposed in estimating location parameters and the corresponding variances. An estimate of bias and a measure of variance of the point estimate are computed using the Monte Carlo method. The bootstrap estimator of the population mean is efficient and consistent in the homogeneous, heterogeneous, and two-segment populations simulated. The design-unbiased approximation of the standard error estimate differs substantially from the bootstrap estimate in severely heterogeneous and positively skewed populations.

Keywords:

Design-unbiased approximation

Heterogeneous

and two-segment populations

Homogeneous

Nonparametric bootstrap

Mathematics Subject Classification:

62D05

62F40

Notes

Note: Values in parentheses are the design-unbiased approximations of the estimated standard errors.

(a) Bootstrap estimates using $k = 100$ replications, $n =$ resample size, at trimmed $N = 257$.

Note: Values in parentheses are the design-unbiased approximations of the estimated standard errors.

(a) Bootstrap estimates using $k = 100$ replications, $n =$ resample size, at trimmed $N = 274$.

Note: Values in parentheses are the design-unbiased approximations of the estimated standard errors.

(a) Bootstrap estimates using $k = 100$ replications, $n =$ resample size, at trimmed $N = 277$.

Note: Resample size = n .

Related research

People also read

Recommended articles

Cited by
240

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG



Taylor & Francis
by informa