

19,932 Views | 170 CrossRef citations to date | 3 Altmetric

Sampling Theory

# Estimation Under Purposive Sampling

Jacqueline M. Guarte & Erniel B. Barrios

Pages 277-284 | Received 01 Sep 2004, Accepted 01 Aug 2005, Published online: 15 Feb 2007

Cite this article <https://doi.org/10.1080/03610910600591610>

Sample our Computer Science Journals

>> [Sign in here](#) to start your access to the latest two volumes for 14 days

- Full Article
- Figures & data
- References
- Citations
- Metrics
- Reprints & Permissions
- Read this article**

## We Care About Your Privacy

We and our 855 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting "I Accept" enables tracking technologies to support the purposes shown under "we and our partners process data to provide," whereas selecting "Reject All" or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the ["privacy preferences"] link on the bottom of the webpage [or the floating icon on the bottom-left of the webpage, if applicable]. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. [Here](#)

We and our partners process data to provide:

I Accept

Reject All

Show Purpose



thin the  
of interest.  
d the  
the point  
ator of the  
eous, and  
the  
severely

Homogeneous

Keyword

Nonparam

Mathem

# Notes

Note: Values in parentheses are the design-unbiased approximations of the estimated standard errors.

(a) Bootstrap estimates using  $k = 100$  replications,  $n =$  resample size, at trimmed  $N = 257$ .

Note: Values in parentheses are the design-unbiased approximations of the estimated standard errors.

(a) Bootstrap estimates using  $k = 100$  replications,  $n =$  resample size, at trimmed  $N = 274$ .

Note: Values in parentheses are the design-unbiased approximations of the estimated standard errors.

(a) Bootstrap estimates using  $k = 100$  replications,  $n =$  resample size, at trimmed  $N = 277$ .

Note: Resample size =  $n$ .

Relate



Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Keep up to date

Register to receive personalised research and resources by email

 Sign me up

- 
- 
- 
- 
- 

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books



Copyright

Accessib

Registered  
5 Howick Pl

or & Francis Group  
orma business

