

89 Views | 0 CrossRef citations to date | 0 Altmetric

Original Articles

Sample Size and the Accuracy of the Generalized Lambda Distribution

George S. Ford & Sarah J. Skinner 






Pages 631-637 | Received 10 Sep 2008, Accepted 31 Oct 2008, Published online: 05 Feb 2009

 Cite this article  <https://doi.org/10.1080/03610910802592846>

Sample our
Computer Science
Journals

>> **Sign in here** to start your access
to the latest two volumes for 14 days



-  Full Article
-  Figures & data
-  References
-  Citations
-  Metrics
-  Reprints & Permissions
- [Read this article](#)
- [Share](#)

Abstract

The Generalized Lambda Distribution is a popular tool for generating random numbers following a wide range of non normal, asymmetric distributions. In this article, we assess the accuracy of the technique in replicating moments across a range of “sample sizes.” We find that accuracy is highly dependent on sample size, particularly with respect to the third and fourth moments. Generally, acceptable accuracy is accomplished with no less than 2,500 observations, but this finding depends critically on the size of the fourth moment.

Keywords:

[Generalized bootstrap](#) [Monte Carlo](#) [Non normal distribution](#) [Non normality](#) [Simulation](#)

Mathematics Subject Classification:

Notes

¹Table values focus on the skewness and kurtosis (with mean 0 and standard deviation 1). The λ_1 and λ_2 parameters determine the location and spread of the data, and can be adjusted to different values without affecting the λ_3 and λ_4 parameters. The tables also consider positive skew, and negative skew is accomplished by flipping the λ_3 and λ_4 values.

²We are grateful to an anonymous referee for these insights.

³The tables in Karian and Dudewicz ([2000](#)) are based on ($m_1 = 0, m_2 = 1$), but changing m_1 to 1.0 requires nothing more than substituting λ_1 from the table with $(\lambda_1 + 1)$. We use $m_1 = 1$ to avoid division by zero.

⁴The random seed is unique to each simulation but identical across each case. Results were computed using other random number generators with little effect on the findings.

Related research

People also read

Recommended articles

Cited by

[Estimating the Parameters of the Generalized Lambda Distribution: Which Method Performs Best? >](#)

Canan G. Corlu et al.

Communications in Statistics - Simulation and Computation

Published online: 27 Jan 2015

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG



Taylor & Francis
by informa