







Q

Home ▶ All Journals ▶ Mathematics, Statistics & Data Science

- Communications in Statistics Simulation and ComputationList of IssuesVolume 38, Issue 3
- Sample Size and the Accuracy of the Gene

Communications in Statistics - Simulation and Computation >

Volume 38, 2009 - Issue 3

82 Views CrossRef citations to date Altmetric

Original Articles

Sample Size and the Accuracy of the Generalized Lambda Distribution

George S. Ford & Sarah J. Skinner

Pages 631-637 | Received 10 Sep 2008, Accepted 31 Oct 2008, Published online: 05 Feb 2009

66 Cite this article https://doi.org/10.1080/03610910802592846



Full Article

Figures & data

References

Metrics

Reprints & Permissions

Read this article

Share

Abstract

The Generalized Lambda Distribution is a popular tool for generating random numbers following a wide range of non normal, asymmetric distributions. In this article, we assess the accuracy of the technique in replicating moments across a range of "sample" sizes." We find that accuracy is highly dependent on sample size, particularly with respect to the third and fourth moments. Generally, acceptable accuracy is accomplished with no less than 2,500 observations, but this finding depends critically on the size of the fourth moment.

Keywords:

Generalized bootstrap

Monte Carlo

Non normal distribution

Non normality

Simulation

Mathematics Subject Classification:

Notes

¹Table values focus on the skewness and kurtosis (with mean 0 and standard deviation 1). The λ_1 and λ_2 parameters determine the location and spread of the data, and can be adjusted to different values without affecting the λ_3 and λ_4 parameters. The tables also consider positive skew, and negative skew is accomplished by flipping the λ_3 and λ_4 values.

²We are grateful to an anonymous referee for these insights.

³The tables in Karian and Dudewicz (2000) are based on (m $_1$ = 0, m $_2$ = 1), but changing m $_1$ to 1.0 requires nothing more than substituting λ_1 from the table with (λ_1 + 1). We use m $_1$ = 1 to avoid division by zero.

⁴The random seed is unique to each simulation but identical across each case. Results were computed using other random number generators with little effect on the findings.



People also read

Recommended articles

Cited by

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up











Accessibility



Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions



Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG