

Journal of the American College of Nutrition >

Volume 21, 2002 - [Issue 6](#)

9,381 Views | 1,522 CrossRef citations to date | 572 Altmetric

Reviews

Omega-3 Fatty Acids in Inflammation and Autoimmune Diseases

Artemis P. Simopoulos , MD, FACN

Pages 495-505 | Received 21 May 2002, Accepted 15 Aug 2002, Published online: 27 Jun 2013

🗨️ Cite this article [🔗 https://doi.org/10.1080/07315724.2002.10719248](https://doi.org/10.1080/07315724.2002.10719248)

Sample our
Food Science & Technology
Journals
>> **Sign in here** to start your access
to the latest two volumes for 14 days

 Full Article  Figures & data  References  Citations  Metrics

 Reprints & Permissions

Read this article

 Share

Abstract

Among the fatty acids, it is the omega-3 polyunsaturated fatty acids (PUFA) which possess the most potent immunomodulatory activities, and among the omega-3 PUFA, those from fish oil—eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA)—are more biologically potent than α -linolenic acid (ALA). Some of the effects of omega-3 PUFA are brought about by modulation of the amount and types of eicosanoids made, and other effects are elicited by eicosanoid-independent mechanisms, including actions upon intracellular signaling pathways, transcription factor activity and gene expression. Animal experiments and clinical intervention studies indicate that omega-3 fatty acids have anti-inflammatory properties and, therefore, might be useful in the management of inflammatory and autoimmune diseases. Coronary heart disease, major depression, aging and cancer are characterized by an increased level of interleukin 1 (IL-1), a

proinflammatory cytokine. Similarly, arthritis, Crohn's disease, ulcerative colitis and lupus erythematosus are autoimmune diseases characterized by a high level of IL-1 and the proinflammatory leukotriene LTB₄ produced by omega-6 fatty acids. There have been a number of clinical trials assessing the benefits of dietary supplementation with fish oils in several inflammatory and autoimmune diseases in humans, including rheumatoid arthritis, Crohn's disease, ulcerative colitis, psoriasis, lupus erythematosus, multiple sclerosis and migraine headaches. Many of the placebo-controlled trials of fish oil in chronic inflammatory diseases reveal significant benefit, including decreased disease activity and a lowered use of anti-inflammatory drugs.

inflammation

cardiovascular disease and major depression autoimmune diseases

IL-1

IL-6

TNF

background diet

omega-6/omega-3 ratio

Key teaching points:

- In Western diets, omega-6 fatty acids are the predominant polyunsaturated fats. The omega-6 and omega-3 fatty acids are metabolically distinct and have opposing physiologic functions.
- Eicosapentaenoic acid (EPA) is released to compete with arachidonic acid (AA) for enzymatic metabolism inducing the production of less inflammatory and chemotactic derivatives.
- Animal and human studies support the hypothesis that omega-3 PUFA suppress cell mediated immune responses.
- In experimental animals and humans, serum PUFA levels predict the response of proinflammatory cytokines to psychologic stress. Imbalance in the omega-6/omega-3 PUFA ratio in major depression may be related to the increased production of proinflammatory cytokines and eicosanoids in that illness.
- The increased omega-6/omega-3 ratio in Western diets most likely contributes to an increased incidence of cardiovascular disease and inflammatory disorders.
- Patients with autoimmune diseases, such as rheumatoid arthritis, inflammatory bowel disease and asthma, usually respond to eicosapentaenoic acid (EPA) and

docosahexaenoic acid (DHA) supplementation by decreasing the elevated levels of cytokines.

Key teaching points:

- In Western diets, omega-6 fatty acids are the predominant polyunsaturated fats. The omega-6 and omega-3 fatty acids are metabolically distinct and have opposing physiologic functions.
 - Eicosapentaenoic acid (EPA) is released to compete with arachidonic acid (AA) for enzymatic metabolism inducing the production of less inflammatory and chemotactic derivatives.
 - Animal and human studies support the hypothesis that omega-3 PUFA suppress cell mediated immune responses.
 - In experimental animals and humans, serum PUFA levels predict the response of proinflammatory cytokines to psychologic stress. Imbalance in the omega-6/omega-3 PUFA ratio in major depression may be related to the increased production of proinflammatory cytokines and eicosanoids in that illness.
 - The increased omega-6/omega-3 ratio in Western diets most likely contributes to an increased incidence of cardiovascular disease and inflammatory disorders.
 - Patients with autoimmune diseases, such as rheumatoid arthritis, inflammatory bowel disease and asthma, usually respond to eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) supplementation by decreasing the elevated levels of cytokines.
-

Related research

People also read

Recommended articles

Cited by
1522

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG



Taylor & Francis
by informa