

Cancer Investigation >

Volume 26, 2008 - [Issue 2](#)

201 | 23 | 3
Views | CrossRef citations to date | Altmetric

ORIGINAL ARTICLE Imaging, Diagnosis, Prognosis

Metastasis-Associated Protein S100A4 and p53 Predict Relapse in Curatively Resected Stage III and IV (M0) Gastric Cancer

Yu Jung Kim, Min A Kim, Seock-Ah Im, Tae Min Kim, Dong-Wan Kim, Han-Kwang Yang,
... [show all](#)

Pages 152-158 | Published online: 11 Jun 2009

🗨️ Cite this article 🔗 <https://doi.org/10.1080/07357900701518909>

Sample our
Medicine, Dentistry, Nursing
& Allied Health Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

📄 Full Article 🖼️ Figures & data 📖 References 🗨️ Citations 📊 Metrics

📄 Reprints & Permissions

Read this article

🔗 Share

Abstract

Purpose: Pathologic stage is the most important predictive factor of relapse in gastric cancer after curative resection. However, patients with the same stage often have different risks of relapse. Here, we investigated whether the expressions of molecular markers can supplement the current staging system in terms of relapse prediction. **Patients and Methods:** One hundred and nine stage III or IV (M0) patients who had received curative gastrectomy followed by adjuvant 5-fluorouracil and cisplatin chemotherapy were included in this study. The expressions of molecular markers including p53, p27, COX-2, HER-2, EGFR, maspin, S100A4, E-cadherin, Sp1, and p97 were analyzed by immunohistochemistry in cancer and paired normal tissues. **Results:**

The overall relapse rate was 58.7%, and pathologic stage was a significant predictive factor of relapse (42% in stage IIIA, 48% in IIIB, 76% in IV, $p = 0.005$). Of the 10 markers examined, p53 and S100A4 were expressed only in tumor tissues, and S100A4 expression was significantly associated with a higher relapse rate (85% vs. 53%, $p = 0.008$). In multivariate analysis including tumor stage, S100A4 and p53 expression were independent predictive factors of relapse (relative risk, 6.98; 95% confidence interval [CI], 1.608-30.342, 3.49; 95% CI, 1.328-9.186, respectively). On comparing patients who expressed S100A4 or p53 with those who expressed neither, relapse rates were 58% vs. 25% in stage III ($p = 0.011$) and 95% vs. 59% in stage IV (M0) ($p = 0.003$). Conclusion: In addition to staging system, the expressions of S100A4 and p53 were significant predictive factors of relapse in gastric cancer after curative resection and adjuvant chemotherapy.

Keywords:

Gastric cancer

Relapse

Immunohistochemistry

S100A4

p53



Related research 

People also read

Recommended articles

Cited by
23

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG

 Taylor and Francis
Group