



Home ▶ All Journals ▶ Medicine ▶ Cancer Investigation ▶ List of Issues ▶ Volume 26, Issue 2 ► Metastasis-Associated Protein S100A4 and ....

Cancer Investigation > Volume 26, 2008 - Issue 2

194 23

Views CrossRef citations to date Altmetric

ORIGINAL ARTICLE Imaging, Diagnosis, Prognosis

## Metastasis-Associated Protein S100A4 and p53 Predict Relapse in Curatively Resected Stage III and IV (M0) Gastric Cancer

Yu Jung Kim, Min A Kim, Seock-Ah Im, Tae Min Kim, Dong-Wan Kim, Han-Kwang Yang,

Pages 152-158 | Published online: 11 Jun 2009

**66** Cite this article ▲ https://doi.org/10.1080/07357900701518909

Sample our Medicine, Dentistry, Nursing & Allied Health journals, sign in here to start your FREE access for 14 days

Full Article

Figures & data

References

**66** Citations

**Metrics** 

Reprints & Permissions

Read this article

Share

## Abstra

Purpose

cancer a

different

markers

**Patients** 

recei chem

including

were an The ove

factor of

markers

expressi

## We Care About Your Privacy

We and our 912 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting I Accept enables tracking technologies to support the purposes shown under we and our partners process data to provide. Selecting Reject All or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the Show Purposes link on the bottom of the webpage . Your choices will have effect within our Website. For more details, refer to our Privacy Policy. Here

We and our partners process data to provide:

Use precise geolocation data. Actively scan device

I Accept in gastric Reject All n have Show Purpose molecular diction. no had atin

> and p97 es. Results: predictive  $e^{10}$


rkers

and S100A4

53%, p =

0.008). In multivariate analysis including tumor stage, S100A4 and p53 expression were independent predictive factors of relapse (relative risk, 6.98; 95% confidence interval [CI], 1.608-30.342, 3.49; 95% CI, 1.328-9.186, respectively). On comparing patients who expressed S100A4 or p53 with those who expressed neither, relapse rates were 58% vs. 25% in stage III (p = 0.011) and 95% vs. 59% in stage IV (M0) (p = 0.003). Conclusion: In addition to staging system, the expressions of S100A4 and p53 were significant predictive factors of relapse in gastric cancer after curative resection and adjuvant chemotherapy.

## Keywords:



Information for Open access Authors Overview R&D professionals Open journals Editors **Open Select** Librarians **Dove Medical Press** Societies F1000Research Opportunities Help and information Reprints and e-prints Advertising solutions Newsroom Accelerated publication Corporate access solutions Books Keep up to date Register to receive personalised research and resources by email Sign me up X or & Francis Group Copyright