







Q

Home ▶ All Journals ▶ Medicine ▶ Gynecological Endocrinology ▶ List of Issues ▶ Volume 34, Issue 2 ▶ Evaluation of pro-oxidant-antioxidant ba

Gynecological Endocrinology >

Volume 34, 2018 - Issue 2

545 97
Views CrossRef citations to date Altmetric
PCOS: PAB and Inflammatory Cytokines

Evaluation of pro-oxidant-antioxidant balance (PAB) and its association with inflammatory cytokines in polycystic ovary syndrome (PCOS)

T. Artimani ☑, J. Karimi, M. Mehdizadeh, M. Yavangi, E. Khanlarzadeh, M. Ghorbani, ...show all

Pages 148-152 | Received 22 Feb 2017, Accepted 22 Aug 2017, Published online: 04 Sep 2017

















Read this article




Abstract

Chronic low-grade inflammation has been suggested as a key contributor of the pathogenesis and development of polycystic ovary syndrome (PCOS). To investigate the association between oxidative stress status and inflammatory cytokines in follicular fluid of 21 PCOS women compared to 21 women with normal ovarian function who underwent intra-cytoplasmic sperm injection. Concentration of IL-6, IL-8, IL-10, and TNF- α was measured using sandwich ELISA. Oxidative stress was examined by measuring total oxidant status (TOS), malondialdehyde (MDA), total antioxidant capacity (TAC), and thiol groups. PCOS women had an elevated concentration of MDA and TOS

compared to controls. Levels of TAC and thiol groups were lower in PCOS compared to controls. PCOS patients had a higher concentration of IL-6, IL-8, and TNF- α compared to controls. Concentration of IL-10 was lower in PCOS compared to controls. Significant correlations were found between MDA and TOS concentration with TNF- α and between IL-6 and MDA, IL-8 and TAC, IL-10 and TOS levels and also between IL-10 and TAC levels. TAC and thiol groups were negatively correlated with TNF- α . Increased oxidative stress in PCOS is associated with inflammation which is closely linked. Inflammation can induce production of inflammatory cytokines in this syndrome and directly stimulates excess ovarian androgen production.

Chinese abstract



Keywords:

Oxidative stress	inflammation	inflammatory cytokines	follicular fluid	PCOS

Acknowledgements

This work was supported by Research and Technology Deputy of Hamadan University of Medical Sciences (IR.UMSHA.REC.1395.191). We express our appreciation to the Research Center of Endometrium and Endometriosis, Hamadan University of Medical Sciences, for their clinical evaluation of patients.

The study was approved by Hamadan University of Medical Science Ethics Committee (IR.UMSHA.REC.1395.191). All subjects provided written informed consent. All procedures performed were in accordance with the ethical standards of the local research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Disclosure statement

The authors report no conflicts of interest.

Additional information

Funding

This work was supported by Research and Technology Deputy of Hamadan University of Medical Sciences. We express our appreciation to the Research Center of Endometrium and Endometriosis, Hamadan University of Medical Sciences, for their clinical evaluation of patients.



People also read

Recommended articles

Cited by 97

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up











Accessibility



Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions



Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG