



Q

Home ► All Journals ► Medicine ► Physiotherapy Theory and Practice ► List of Issues ► Volume 32, Issue 7 ► Kinesio taping or just taping in shoulde ....

Physiotherapy Theory and Practice >

An International Journal of Physical Therapy

Volume 32, 2016 - <u>Issue 7</u>

1,741177ViewsCrossRef citations to dateAltmetricResearch ReportResearch ReportAltmetric

# Kinesio taping or just taping in shoulder subacromial impingement syndrome? A randomized, double-blind, placebo-controlled trial

Figen Kocyigit S, MD, Merve Acar, MD, Mehmet Besir Turkmen, MD, Tugce Kose, PT, Nezahat Guldane, PT & Ersin Kuyucu, MD

Pages 501-508 | Received 10 Mar 2015, Accepted 11 Oct 2015, Published online: 12 Sep 2016

<b>66</b> Cite this article	https:/	/doi.org/10	.1080/095939	R Check for updates		
				entistry, Nursing		
			& Allied Heal >> Sign in here to the latest two	th Journals to start your access volumes for 14 days		
Full Article	🖾 Figures	& data	Reference	es <b>66</b> Citations	s 🔟 Metrics	
A Reprints & Permissions Read the Read the Reprints & Read the Reprint Read the Reprint Read the Reprint Read the Reprint Repr		is article	Share			

## ABSTRACT

Objective: To verify effects of kinesio taping (KT) in shoulder subacromial impingement syndrome (SIS) when compared to sham taping applied in the same way with KT. Patients and Methods: Patients were randomized as group 1 (n = 21) KT group and group 2 (n = 20) sham-taping group. Taping was applied every three days, three times during the study period. We assessed all the patients at baseline, at the end of the taping period (12th day), and at one-month post-intervention. We assessed pain on the 100 mm visual analog scale (VAS). Shoulder range of motion (ROM), Constant Scores, and Nottingham Health Profile (NHP) scores were evaluated. Results: Of the 41 participants, 13 were males (32%) and 28 were females (68%). The mean age was 45 ±

15 years (range 20–65 years). We documented a significant decrease in VAS for nocturnal pain, and Constant Score in both groups. The KT group showed additional significant change in NHP pain and physical activity scores. Conclusion: KT and sham taping generated similar results regarding pain and Constant Scores.

**KEYWORDS**:



### Declaration of interest

The authors declare that they have no conflict of interest.

#### Related research 1

People also read Recommended articles	Cited by 17
---------------------------------------	----------------

Information for	Open access
Authors	Overview
R&D professionals	Open journals
Editors	Open Select
Librarians	Dove Medical Press
Societies	F1000Research
Opportunities	Help and information
Reprints and e-prints	Help and contact
Advertising solutions	Newsroom
Accelerated publication	All journals
Corporate access solutions	Books

#### Keep up to date

Register to receive personalised research and resources by email





Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions

Taylor & Francis Group an informa business



Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG