



Q

Home ► All Journals ► Medicine ► Stress ► List of Issues ► Volume 20, Issue 1 ► Toll-like receptor 4 protects against st

Stress >

The International Journal on the Biology of Stress Volume 20, 2017 - <u>Issue 1</u>

✓ Free access

766
Views2
CrossRef citations to date0
Altmetric

Listen

Original Research Report

Toll-like receptor 4 protects against stressinduced ulcers via regulation of glucocorticoid production in mice

Liang Wang, Pengfei Luo, Fang Zhang, Yuelu Zhang, Xingtong Wang, Fei Chang, ... show all Pages 19-26 | Received 10 Sep 2015, Accepted 11 Aug 2016, Published online: 06 Dec 2016

L Cite this article **I** https://doi.org/10.1080/10253890.2016.1224843

Check for updates

I Accept

🖹 Full Ar

🔒 Repri

Abstra

Stress-ir molecula (TLR4), a trans prote (WT) and and exp post mo mice. Fu observed

We Care About Your Privacy

We and our 912 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting "I Accept" enables tracking technologies to support the purposes shown under "we and our partners process data to provide," whereas selecting "Reject All" or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the ["privacy preferences"] link on the bottom of the webpage [or the floating icon on the bottom-left of the webpage, if applicable]. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. <u>Here</u>

We and our partners process data to provide:

. -

Reject All Show Purpose the eptor 4 immatory played a 5. Wild-type

> er group), determined FLR4^{–/–} ons were

ce 4 h after

eover. the

. .

showed expression in the adrenal of P450scc (CYP11A1), the first rate-limiting enzyme in the synthesis of steroids, was increased 4 h after water immersion restraint stress or LPS treatment in WT mice, but was conversely decreased in TLR4^{-/-} mice after either stressor. Furthermore, in adrenal glands of TLR4^{-/-} mice, structural distortion of mitochondria (which contain CYP11A1) was found with electron microscopy, and lack of lipid-storing droplets was found using light microscopy on adrenal cryosections stained with Oil red O. These data indicate that TLR4 plays a protective role in stress-induced gastric ulcer that is exerted via impacting synthesis of glucocorticoid in the adrenal gland.

Keywords:

Cortisol	alucocorticoid	I PS	stomach ulcer	stress	TIR4
COLISOI	giucocorticolu	LFJ	Stomach uicei	SUESS	ILN4

Introduction

Stress may affect different physiological functions of the gastrointestinal tract including gastric secretion, motility, permeability and barrier function, visceral sensitivity, and

mucosal		ould lead to
gastric u	×	er is a
commor		(Ali & Harty,
<u>2009</u>). Т		d not fully
understo		ts starting
with the		ulceration by
physiolo		echolamines
and provide the second se		
Endo		during acute
stress ha		aretova
et al., <u>20</u>		, activated
in respon		ropin-
releasing		5
gastropr		<u>8, 2012</u>).
Moreove		age
Article contents	🛓 Related research	

Moreover, intra-hypothalamic dexamethasone implantation which induces long-lasting stress-induced glucocorticoid deficiency results in aggravation of stress-induced gastric injury (Filaretova & Filaretov, 1992). The gastroprotective effects of glucocorticoids during acute stress involve multiple mechanisms, including the maintenance of glucose homeostasis, gastric mucosal blood flow, mucus production, and attenuation of gastric motility and microvascular permeability (Filaretova et al., 2004; Kuo et al., 2015; Podvigina et al., 2009), as well as changes in the gastric microcirculation (Filaretova et al., 1999). Although circulating corticosterone is considered to be the main glucocorticoid in the regulation of stress in rodents, cortisol also has a corresponding response to stimulation by stress in mice (Gong et al., 2015). Whether it impacts stress-induced gastric ulcer in mice remains to be determined.

TLR4 is an innate immune pattern recognition receptor, which is part of the Interleukin-1 Receptor/Toll-like Receptor Superfamily containing a toll-like/IL-1 Receptor (TIR) domain and a Leucine-rich repeat motif in the extracellular domain (Nardone & Compare, <u>2014</u>). As the receptor for lipopolysaccharide (LPS), TLR4 can be activated by the LPS-lipopolysaccharide-binding protein (LBP)-myeloid differentiation protein 2 (MD2) complex, thus passing this signal to the downstream nuclear factor (NF)-kb pathway (Akira & Takeda, <u>2004</u>). Several studies have reported that downregulation of NF-kb or inflammation mitigates gastric ulcer (Kang et al., <u>2014</u>; Mahmoud-Awny et al., <u>2015</u>;

Sinha et		oromote
stress-in	×	activity.
However		e HPA axis
(Monroe		of CRF (CRH)
gene exi		oum-Ribot
et al., <u>2(</u>		stress-
induced		cocorticoid.
Take		ne TLR4 and e TLR4
downstre		luced gastric
damage		also could
enhance		ological
ulceratic		hesis of
glucocor		ere
overning		
Article contents	🚔 Related research	

Methods

Animals and management

The study was approved by the Institutional Animal Care and Use Committee (IACUC) of the Second Military Medical University in Shanghai (IACUC-2012-XuSG) in accordance with the U.S.A. National Institutes of Health guidelines for the care and use of laboratory mice. TLR4 knockout (TLR4^{-/-}; C57BL/10ScNJU) and wild-type (WT;C57BL/6_129) mice were obtained from the Jackson Laboratory (Bar Harbor, ME, U.S.A.). The TLR4^{-/-} mice were genetically identified in the Nanjing Biomedical Research Institute of Nanjing University (Nanjing, Jiangsu, China). Male WT and TLR4^{-/-} mice bred under specific pathogen-free conditions aged 6-8 weeks were used for experiments. Mice were housed three to five per cage and acclimatized to standard laboratory conditions (lights on between 08:00 h and 19:00 h, temperature 20 ± 1 °r, free access to food and water) for 7 days before the experiment.

Induction of ulcers by exposure to combined water immersion and restraint stress



WT and TLR4^{-/-} mice were fasted for 24 h before experiments but given free access to

water immersion restraint stress (Jia et al., <u>2007b</u>), mice were anesthetized with sodium pentobarbital (as above) for blood sampling and stomach removal for the estimation of cortisol level and gastric injury, respectively.

Endotoxemia model

WT and TLR4^{-/-} mice were randomized (n = 5 per group) and injected intraperitoneally with LPS (0.2 mg/ml, 2 mg/kg) (Sigma-Aldrich Corp., St. Louis, MO) or isotonic saline and killed 4 h later to determine the effect of TLR4 (receptor for LPS) on the regulation of endogenous glucocorticoids.

Corticotropin (adrenocorticotropic hormone, ACTH) test

WT and TLR4^{-/-} mice were randomized and injected intraperitoneally with corticotropin (5 IU/ml, 50 IU/kg) (Shanghai NO.1 Biochemical & Pharmaceutical Co., LTD, Shanghai, China) or isotonic saline and killed after 4 h to measure serum cortisol and corticosterone.

Assessment of gastric damage

Stomachs were removed and filled with 2 ml of 1% formalin and immersed in formalin for 24 h. The stomachs then were cut along the length of the greater curvature and

examine	×	erosions
was mea		tor blinded
to the tr		tissue
sample		
Measu		
lipopol		
Samp		parator tube
and		rations in
serum sa		nl;
coefficie		9.3%-
21.2%) (ssay Kit
(sensitiv		5%, and
inter-ass		Р
concenti		-linked
Article contents	Related research	

<8%, and inter-assay <10%) (Cusabio, Wuhan, China) and the mouse LBP enzymelinked immunosorbent assay kits (sensitivity: 3.12 ng/ml; coefficient of variation: intraassay <8%, and inter-assay <10%) (Cusabio, Wuhan, China).

Western blotting

Whole adrenal glands were lysed in protein extraction buffer [20 mM Tris (pH7.5), 150 mM NaCl, 1%Triton X-100, 1 mM EDTA] followed by centrifugation at 12,000 × g for 15 min at 4 °C. The supernatant was removed and total protein was determined (Bradford assay). Samples containing 40 µg protein were loaded, separated by 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (90 V), and blotted onto nitrocellulose membrane. The blots were probed with anti-CYP11A1 antibody (1/1000; Cell Signaling Technology) and with horseradish peroxidase-conjugated anti-rabbit secondary antibody (1/2000; Santa Cruz Biotechnology). After extensive washing, protein bands were visualized by chemiluminescence staining. Blots were also probed with anti-glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (1/1000; Sigma-Aldrich) to confirm equal loading of protein. Expression of proteins was then calculated semiquantitatively as integrated optical density (IOD) ratio of CYP11A1 to GAPDH.

Electron microscopy



Statistical analysis

Data were analyzed by two-way ANOVA, one-way ANOVA, or Tukey's test of variance using SPSS v.20.0 software (SPSS Inc., Chicago, IL). Differences between groups were calculated using the least significant differences method, and significance level was defined as p < .05. Results are expressed as mean \pm SEM.

Results

TLR4 deficiency exacerbates stress-induced gastric mucosal injury and modulates serum glucocorticoid concentrations

Gastric mucosa. Gastric mucosal injury was exacerbated in WT mice as a function of water immersion restraint time. Cellular swelling, hypochromia of the cytoplasm, stenosis, and erythropenia in the mucosa appeared starting 0.5 h after stress application. Atrophy and exfoliation of mucosal cells, and increased gastric stromal cell erosion was observed at 1 h. After 2 h, mucosal erosion became more severe, and ulceration appeared at 4 h. The pathological changes induced by water immersion restraint stress were more severe in TLR4^{-/-} than in WT mice (Figure 1(A)), and more ulcers were observed grossly in the gastric mucosa of the mutants after 4 h (Figure 1(B))

), which	X	ANOVA,
F _{mice-type}		=26.8,
Pmice-type		= 123.7, p
<.001;		Г1h/TLR4—/—1 h
<.001, p		
Figure 1		(A)
Histol) exposed
for in		of mice (n =
5 each)		ied as
gastric u		sterone
concenti		05 (one-way
ANOVA,		



Serum cortisol. Serum cortisol concentrations were increased in WT mice starting from 2 h after the application of the stressor and peaked at the 4-h time point (two-way ANOVA, $F_{\text{mice-type}} = 109.4$, $p_{\text{mice-type}} < .001$; $F_{\text{time}} = 29.8$, $p_{\text{time}} < .001$; $F_{\text{mice-type} \times \text{time}} =$ 30.7, $p_{mice-type \times time} < .001$; one-way ANOVA after two-way ANOVA, df = (9,40), F = 39.0, p < .001; Tukey's test, $p_{WT0h/WT0.5h} = 1.000$, $p_{WT0.5h/WT1h} = .39$, $p_{WT1h/WT2h} = .001$, $p_{WT2h/WT4h} = .005$). In contrast, cortisol concentration remained lower in TLR4^{-/-} mice during exposure to stress (one-way ANOVA after two-way ANOVA, df = (9, 40), F = 39.0, p < .001; Tukey's test, $p_{WT0h/TIR4-/-0h} = .8$, $p_{WT0.5h/TIR4-/-0.5h} = .8$, $p_{WT1h/TIR4-/-1h} = .3$, PWT2h/TL × by the Serum c pes of mice stressor (two-wa F_{mice-type ×} = (9,40), F $t_{ime} = 62$ = 75.7, $_{\rm WT2h}$ < .001, $O_{TLR4-/-1}$ PWT2h/ ed a higher h/WT2 basal va e to the 001; Tukey's stressor test, pw⁻)1, p_{WT 2} h/TLR4-/osal injury Admini

As expected, serum cortisol concentrations were increased in WT and TLR4^{-/-} mice upon injection of exogenous cortisol as compared to the vehicle-injected and uninjected control groups (WT: F = 2.98, df = (2, 12) p = .0001; TLR4^{-/-}: F = 67.18, df = (2, 12), p < .0001) (Figure 2(A)) and was associated with a lower UI (WT: F = 19.02, df = (2, 12) p = .0002; TLR4^{-/-}: F = 60.13, df = (2, 12), p < .0001) (Figure 2(B)). Hence, cortisol administration alleviated stress-induced gastric mucosal damage, in both WT and TLR4^{-/-} mice.

Figure 2. Effect of cortisol administration on stress-induced ulceration. WT and TLR4^{-/-} mice were evaluated for (A) serum cortisol concentrations, and (B) gastric ulcer index (UI) after pretreatment with cortisol. Results represent mean ± SEM (n = 5 mice per group). *p < .05 (one-way ANOVA, Tukey's test) versus vehicle group (dimethylsulfoxide in saline, 1:200) and control groups (without any treatment).



Display full size





Cortisol and corticosterone concentrations were measured in mice pretreated with LPS (1% w/v, 2 mg/kg) for 4 h. Serum LPS concentration increased compared to the vehicle-treated group in both WT and TLR4^{-/-} mice (WT: F = 123.4, p < .0001; TLR4^{-/-}: F = 35.14, p = .0004. All df are (3, 16)), while cortisol concentrations were elevated in LPS-treated as compared to vehicle-treated WT but not TLR4^{-/-} mice (WT: F = 33.03, p



microscopy, obvious reduction in density of lipid-storing droplets was seen in Oil red Ostained whole adrenal frozen section in unstressed TLR4^{-/-} mice (Figure 4(B)).

Figure 4. Ultrastructure, lipid storage, and steroidogenic enzyme content in WT and TLR4^{-/-} mouse adrenal glands. (A) Electron micrographs of adrenocortical cells in the zona fasciculata. Mitochondria (MIT), smooth endoplasmic reticulum, and liposomes (LIP) are visible in the cytoplasm of WT and TLR4^{-/-} mice. Note the round shape, disorganized arrangement of tubulovesicular cristae, and presence of metrical vacuoles in the mutant. Scale bar 1 μ m. (B) Cryosections of whole adrenal gland stained with Oil red O in WT and TLR4^{-/-} mice without any treatment showing fewer red-stained lipid-droplets in TLR4 mutant mice. Scale bar 200 μ m. (C) Western blotting of adrenal gland protein samples and the corresponding integrated optical density (IOD) ratio values (CYP11A1/glyceraldehyde 3-phosphate dehydrogenase [GAPDH]). The expression of CYP11A1 (P450scc) increased 4 h after stress or lipopolysaccharide (LPS) treatment in WT but decreased in TLR4^{-/-} mice. Results represent mean ± SEM (n = 5 mice per group). *p < .05 versus WT vehicle group (isotonic saline). #p < .05 (one-way ANOVA, Tukey's test) versus TLR4^{-/-} vehicle group (isotonic saline).



Discussion

The activation of the inflammatory response and ensuing release of pro-inflammatory cytokines are features of stress-induced peptic ulcers (McGettrick & O'Neill, 2004). The innate inflammatory/immune response is activated after exposure to stressful stimuli. Cold restraint stress can induce a rapid activation of nuclear factor (NF)-kB in rats, the inhibition of which alleviates inflammation and injury (Jia et al., <u>2007a</u>). The deletion of the NF-κB p50 subunit can alleviate gastric injury by inhibiting NF-κB activation and consequent inflammation (Ye et al., 2013). We previously demonstrated that the activation of the mitogen-activated protein kinase (MAPK) p38 by reactive oxygen species may also play a role in stress-induced gastric damage in rats (Jia et al., 2007b). TLR4 protein is a positive regulator of inflammation (Szumilas et al., 2013). Signaling pathways activated by Toll-like receptors (TLRs) in the innate immune system are linked to the induction of pro-inflammatory cytokines. NF-κB, p38, and the MAPK family member c-Jun N-terminal kinase are the major factors activated by TLR4 (O'Neill, 2006) which, along with its co-receptor MD-2 and some inflammatory mediators, is upregulated by chronic mild stress (Garate et al., 2011). Hence it is a plausible hypothesis that TLR4 may increase gastric injury through the activation of inflammatory pathways in stress ulcer. However, we found just the opposite. In this study, it was observed that water immersion restraint stress-induced gastric ulceration



WT and TLR4^{-/-} mice. This indicates that the release of glucocorticoid during acute stress plays an important role in indicating injury and protecting the gastric mucosa during acute stress in mice.

Previous research showed that TLR4 signaling is sufficient to cause glucocorticoid release from adrenal cells (Vakharia & Hinson, 2005). LPS from Gram-negative bacteria, which act via TLR4, stimulates glucocorticoid release (Zacharowski et al., 2006). In addition, translocation from the intestinal flora of bacterial LPS occurs during chronic mild stress (Garate et al., 2011). LBP is essential for the recognition of LPS by TLR4 (Palsson-McDermott & O'Neill, 2004), and it has been found that plasma LPS and LBP expression is upregulated after cold restraint stress, which has been linked to changes in intestinal permeability and bacterial translocation (Garate et al., 2011). In this study, we also found that serum concentrations of LPS and of LBP were upregulated after 4-h water immersion restraint stress. Also, the serum cortisol and corticosterone concentrations both increased 4 h after LPS treatment in this study in WT, but importantly not in the TLR4^{-/-} mice. In contrast, the corticotropin test showed increased circulating glucocorticoid (corticosterone and cortisol) concentrations in both types of mice, although the response was less in $TLR4^{-/-}$ mice. Hence, TLR4 in the adrenal gland is not essential for a response to ACTH, but has an important role in regulating the extent of glucocorticoid release during acute stress.

Convers	×	limiting, and
hormona		on et al.,
<u>1961;</u> Ka		sis is
controlle		h
conversi		leavage
enzyme		ace in the
endopla		CYP11A1
engi		oplasmic
reticu		s required
for activ		ocortical
cells of 7		of WT mice.
Furthern		nd by LPS
treatme		storing
droplets		R4 ^{-/-} mice
in Oil roo		wunderlie
Article contents	📥 Related research	

the differences in the synthesis and secretion of corticosterone and cortisol by adrenocortical cells between WT and $TLR4^{-/-}$ mice.

Exposure to low concentrations of endotoxin produces long-term changes in HPA axis activity (Shanks et al., 2000). In particular, early-life exposure to Gram-negative LPS increases corticosterone secretory pulse frequency and amplitude (Vakharia & Hinson, 2005). Moreover, preconditioning with mild stress provides gastroprotection to rats during acute stress (Filaretova et al., 2008). In another study, increasing glucocorticoid receptor expression induced a weaker response to restraint stress (Reichardt et al., 2000). Thus, it can be conjectured that long-term exposure to stressful environmental stimuli may promote the neuroendocrine/neuroimmune response during acute stress events, and that this form of adaptation is lost in the absence of TLR4, thereby decreasing protection against the adverse consequences of acute stress.

In summary, TLR4 deficiency resulted in aggravated gastric ulceration that was associated with a reduction in glucocorticoid release during acute stress, an effect that was alleviated by treatment with exogenous cortisol. TLR4 thus plays a gastroprotective role in stress-induced gastric ulceration that is exerted via modulation of glucocorticoid secretion by impacting the ultrastructure and the synthetic function of the adrenal gland through LPS/TLR4 signaling. In addition, as well as corticosterone, cortisol also plays an important role during acute stress in mice

	×	
Disclos		
No confl		
Refer		
1. Aguile		althy aging.
Exp G		

2. Akira S, Takeda K. (2004). Toll-like receptor signalling. Nat Rev Immunol 4:499–511.

PubMed Web of Science ® Google Scholar

3. Ali T, Harty RF. (2009). Stress-induced ulcer bleeding in critically ill patients. Gastroenterol Clin North Am 38:245–65.

PubMed Web of Science ® Google Scholar

 Black SM, Harikrishna JA, Szklarz GD, Miller WL. (1994). The mitochondrial environment is required for activity of the cholesterol side-chain cleavage enzyme, cytochrome P450scc. Proc Natl Acad Sci USA 91:7247–51.



PubMed Web of Science @ Google Schola

 Filaretova L, Bagaeva T, Morozova O. (2012). Stress and the stomach: corticotropinreleasing factor may protect the gastric mucosa in stress through involvement of glucocorticoids. Cell Mol Neurobiol. 32:829–36.

6. Filaretova I Maltac	N Deaderey A Le	v/covieb \/ (1000)	Dala of apotric	1
micro			×	g water-
restra				
PubMe				
7. Filaret				tective to
ulcero				ion. Curr
Pha				
8. Filaret				f
glucod				nst stress-
induce				
No.				
Article co	ontents		Related research	

 Filaretova LP, Filaretov AA. (1992). [The effect of large doses of corticosteroids on stomach ulcer formation: a new hypothesis] [Article in Russian]. Fiziol Zh SSSR Im I M Sechenova 78:77–83.

PubMed Google Scholar

.0. Filaretova LP, Filaretov AA, Makara GB. (1998). Corticosterone increase inhibits stressinduced gastric erosions in rats. Am J Physiol 274:G1024–30.

Google Scholar

1. Filaretova LP, Podvigina TT, Bagaeva TR, Tanaka A, Takeuchi K. (2004). Mechanisms underlying the gastroprotective action of glucocorticoids released in response to ulcerogenic stress factors. Ann N Y Acad Sci 1018:288–92.

PubMed Web of Science ® Google Scholar

.2. Garate I, Garcia-Bueno B, Madrigal JL, Bravo L, Berrocoso E, Caso JR, Mico JA, Leza JC. (2011). Origin and consequences of brain Toll-like receptor 4 pathway stimulation in an experimental model of depression. J Neuroinflammation 8:151. PubMed PMID:22053929.



PubMed	Web of Science ®	Google Scholar

.6. Jia YT, Ma B, Wei W, Xu Y, Wang Y, Tang HT, Xia ZF. (2007a). Sustained activation of nuclear factor-kappaB by reactive oxygen species is involved in the pathogenesis of stress-induced gastric damage in rats. Crit Care Med 35:1582–91.

PubMed Web of Science ® Google Scholar

.7. Jia YT, Wei W, Ma B, Xu Y, Liu WJ, Wang Y, Lv KY, et al. (2007b). Activation of p38 MAPK by reactive oxygen species is essential in a rat model of stress-induced gastric mucosal injury. J Immunol 179:7808–19.

PubMed Web of Science ® Google Scholar

.8. Kang JW, Yun N, Han HJ, Kim JY, Kim JY, Lee SM. (2014). Protective effect of flos lonicerae against experimental gastric ulcers in rats: mechanisms of antioxidant and anti-inflammatory action. Evid Based Complement Alternat Med 2014:596920.

PubMed Web of Science ® Google Scholar

.9.	Koritz SB, Kumar AM. (1970). On the me	echanism of action of the	e adrenocorticotrophic
	hormc		× olone
	synthe		
0	Κυο Τ		postasis hy
. 0.			.03(03)3 0 9
	gracot		
21.	Loum-		coids down-
	regula		nRNA in the
	rat hy		ons.
	Neuro		
	Article contents	- Polatod	racaarch

 Mahmoud-Awny M, Attia AS, Abd-Ellah MF, El-Abhar HS. (2015). Mangiferin mitigates gastric ulcer in ischemia/reperfused rats: involvement of PPAR-γ, NF-κB and Nrf2/HO-1 signaling pathways. PLoS One 10:e0132497.

Google Scholar

23. McGettrick AF, O'Neill LA. (2004). The expanding family of MyD88-like adaptors in Toll-like receptor signal transduction. Mol Immunol 41:577–82.



 Miller WL, Auchus RJ. (2011). The molecular biology, biochemistry, and physiology of human steroidogenesis and its disorders. Endocr Rev 32:81–151.

PubMed Web of Science ® Google Scholar

 Monroe SM, Simons AD. (1991). Diathesis-stress theories in the context of life stress research: implications for the depressive disorders. Psychol Bull 110:406–25.

6. Nardore C. Compare D. (2014). The neurohe and costric functions. Dir D. X	5 32:206-12.
27. O'Neil know.	we don't
8. Pa	
9. Podviç	etova LP.
(2009	ons on the
Article contents	d hormone

PubMed Google Scholar

30. Quenot JP, Thiery N, Barbar S. (2009). When should stress ulcer prophylaxis be used in the ICU? Curr Opin Crit Care 15:139–43.

PubMed Web of Science ® Google Scholar

31. Reichardt HM, Umland T, Bauer A, Kretz O, Schutz G. (2000). Mice with an increased glucocorticoid receptor gene dosage show enhanced resistance to stress and endotoxic shock. Mol Cell Biol 20:9009–17.



32. Shanks N, Windle RJ, Perks PA, Harbuz MS, Jessop DS, Ingram CD, Lightman SL. (2000). Early-life exposure to endotoxin alters hypothalamic-pituitary-adrenal function and predisposition to inflammation. Proc Natl Acad Sci USA 97:5645–50.

PubMed Web of Science ® Google Scholar

33. Shimozawa N, Okajima K, Harada N, Arai M, Ishida Y, Shimada S, Kurihara H, Nakagata N. (2006). Contribution of sensory neurons to sex difference in the develo erology X 131:1 34. Sinha ic mucosa from r atory iys Acta dam drenals: 5. Stone h Biochem aspec Biophy Article contents

36. Szumilas D, Krysiak R, Okopien B. (2013). [The role of TLR4 receptor in development of inflammation and carcinogenesis in ulcerative colitis and pharmacotherapy of this disorder]. Wiad Lek 66:3–9.

PubMed Google Scholar

 Vakharia K, Hinson JP. (2005). Lipopolysaccharide directly stimulates cortisol secretion by human adrenal cells by a cyclooxygenase-dependent mechanism. Endocrinology 146:1398–402.



 Ye B, Zhou PY, Jia M, Cheng XS, Jia YT, Xu SG. (2013). Absence of NF-kappaB subunit p50 ameliorates cold immobilization stress-induced gastric ulcers. Biochem Biophys Res Commun 434:547–51.

```
PubMed Web of Science ® Google Scholar
```

39. Yigiter M, Albayrak Y, Polat B, Suleyman B, Salman AB, Suleyman H. (2010). Influence of adrenal hormones in the occurrence and prevention of stress ulcers. J Pediatr Surg 45:2154–9.



Information for	Open access
Authors	Overview
R&D professionals	Open journals
Editors	Open Select
Librarians	Dove Medical Press
Societies	F1000Research
Opportunities	Help and information
Reprints and e-prints	Help and contact
Advertising solutions	Newsroom
Accelerated publication	All journals
Corporate access solutions	Books

Keep up to date

Register to receive personalised research and resources by email

ڬ Sign me u

