



Leukemia & Lymphoma >

Volume 49, 2008 - [Issue 11](#)

596 | 51

Views | CrossRef citations to date | 6

Altmetric

Original Articles: Clinical

Hydroxyurea, azacitidine and gemtuzumab ozogamicin therapy in patients with previously untreated non-M3 acute myeloid leukemia and high-risk myelodysplastic syndromes in the elderly: results from a pilot trial

Sucha Nand , John Godwin, Scott Smith, Kevin Barton, Laura Michaelis, Serhan Alkan, ... [show all](#)

Pages 2141-2147 | Received 16 Jun 2008, Accepted 27 Aug 2008, Published online: 01 Jul 2009

Cite this article <https://doi.org/10.1080/10428190802451254>

Sample our
Medicine, Dentistry, Nursing
& Allied Health Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

Read this article

Share

Abstract

Elderly patients with acute myeloid leukemia (AML) and high-risk myelodysplastic syndromes (MDS) have a poor prognosis due to low response rates (26–46%) to standard chemotherapy and high treatment-related mortality (11–31%). In this Phase II study, we used a combination of hydroxyurea (HU), azacitidine and low dose gemtuzumab ozogamicin (GO) to assess its efficacy and toxicity in this group of patients. Twenty patients with non-M3 AML and MDS were treated with this regimen. The treatment was begun with HU 1500 mg orally twice daily to lower white blood cell

count below 10,000/ μ L, followed by azacitidine 75 mg/m² subcutaneously for 7 days and GO 3 mg/m² on day 8. Patients who achieved complete remission (CR) received a consolidation course. The median age of patients was 76 years. Eleven patients (55%) were treated in the outpatient setting. Fourteen (70%) achieved a CR, three of which were incomplete (CRi). The median duration of remission was 8 months and median survival was 10 months. Performance status of 0–1 was associated with high complete response rate. Overall toxicity was acceptable with only one (5%) early death due to disease progression. The combination of HU, azacitidine and GO appears to be a safe and effective regimen in the treatment of AML and high risk MDS in the elderly. These results need to be confirmed in a larger cohort of patients.

Keywords:

- Acute myeloid leukemia
- azacitidine
- gemtuzumab ozogamicin

Related research

People also read

Recommended articles

Cited by
51

Gemtuzumab ozogamicin (GO) as monotherapy and in combination with hypomethylation agents in the treatment of acute myeloid leukemia: a systematic meta-analysis >

Jun Li et al.
Hematology
Published online: 27 Feb 2022



Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)

 Taylor and Francis Group

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG