



Journal of Applied Aquaculture >

Volume 20, 2008 - [Issue 1](#)

267 Views | 12 CrossRef citations to date | 0 Altmetric

Original Articles

Productivity and Economics of Nile Tilapia *Oreochromis niloticus* Cage Culture in South-East Brazil

L. Conte, D.Y. Sonoda, R. Shirota & J.E.P. Cyrino

Pages 18-37 | Published online: 11 Oct 2008

Cite this article <https://doi.org/10.1080/10454430802022060>

Sample our
Environment & Agriculture
Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

Read this article

Share

ABSTRACT

Fish cage culture is an intensive, continuous-flow fish farming system, allowing intensive exploitation of water bodies with relatively low capital investment. This study aimed to determine the production function of Nile tilapia, *Oreochromis niloticus*, in cages; the profit-maximizing biomass at 300–400 and 500–600 fish per m³ for cages of different volumes; and the influence of water body conditions in fish performance. Feed intake, survival rate, and water temperature were monitored daily; dissolved oxygen, pH, and transparency of water were monitored each 15 days. Caged tilapia were fed daily on commercial, floating pellets (32% crude protein) at 0900, 1300, and 1700, and feeding rate was adjusted based on weight gain and survival rate. Data were analyzed statistically by ANOVA ($P = 0.05$) and regression analysis; the Mitscherlich function was chosen to represent the production function. Carrying capacity of both stocking densities reached 200 kg/m³ and no differences were found ($P > 0.05$) regarding

accumulated biomass and individual average weight over time. The larger stocking density yielded larger accumulated biomass and had better feeding efficiency and no differences between individual average weights of fish at both densities were observed ($P > 0.05$). Profit-maximizing biomass at 500–600 fish/m³ was 145 kg/m³ and at 300–400 fish/m³ was 121 kg/m³. Cage farming of Nile tilapia at 500–600 fish/m³, individual average weight 283 g, presented many advantages: optimization of space and production time, better feed efficiency, higher fish production per unit volume of cages, and increased profitability.

KEYWORDS:

- Oreochromis niloticus
- cage farming
- production functions
- biomass

Related research ⓘ

- People also read
- Recommended articles
- Cited by 12

Information for

Authors
R&D professionals
Editors
Librarians
Societies

Opportunities

Reprints and e-prints
Advertising solutions
Accelerated publication
Corporate access solutions

Open access

Overview
Open journals
Open Select
Dove Medical Press
F1000Research

Help and information

Help and contact
Newsroom
All journals
Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)

 **Taylor & Francis Group**
an informa business

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG