



Structural Equation Modeling: A Multidisciplinary Journal >

Volume 20, 2013 - Issue 3

554 | 15 | 0  
Views | CrossRef citations to date | Altmetric

Original Articles

# Investigating Factorial Invariance of Latent Variables Across Populations When Manifest Variables Are Missing Completely

Keith F. Widaman, Kevin J. Grimm, Dawnté R. Early, Richard W. Robins & Rand D. Conger  
Pages 384-408 | Published online: 22 Jul 2013

Cite this article <https://doi.org/10.1080/10705511.2013.797819>

Sample our  
Mathematics & Statistics  
Journals  
**>> Sign in here to start your access  
to the latest two volumes for 14 days**

Full Article Figures & data References Citations Metrics

Reprints & Permissions

Read this article

Share

## Abstract

Difficulties arise in multiple-group evaluations of factorial invariance if particular manifest variables are missing completely in certain groups. Ad hoc analytic alternatives can be used in such situations (e.g., deleting manifest variables), but some common approaches, such as multiple imputation, are not viable. At least 3 solutions to this problem are viable: analyzing differing sets of variables across groups, using pattern mixture approaches, and a new method using random number generation. The latter solution, proposed in this article, is to generate pseudo-random normal deviates for all observations for manifest variables that are missing completely in a given sample and then to specify multiple-group models in a way that respects the random

nature of these values. An empirical example is presented in detail comparing the 3 approaches. The proposed solution can enable quantitative comparisons at the latent variable level between groups using programs that require the same number of manifest variables in each group.

Keywords:

confirmatory factor analysis

factorial invariance

missing data

structural equation modeling

## Notes

<sup>1</sup>EQS ([Bentler, 2006](#)) scripts and output files for all models presented in this article are available on request from the first author.

<sup>2</sup>Mplus ([Muthén & Muthén, 2007](#)) scripts and output files for all pattern mixture models presented in this article are available on request from the first author.

<sup>3</sup>Analysis scripts and output files for all models presented in this article using the random data approach are available on request from the first author.

### Related Research Data

[Maximum Likelihood from Incomplete Data Via the EM Algorithm](#)

Source: Journal of the Royal Statistical Society Series B (Statistical Methodology)

[Integrative data analysis through coordination of measurement and analysis protocol across independent longitudinal studies.](#)

Source: Psychological Methods

[On structural equation modeling with data that are not missing completely at random](#)

Source: Psychometrika

[Exploring the measurement invariance of psychological instruments: Applications in the substance use domain.](#)

Source: Unknown Repository

[Inference and Missing Data](#)

Source: Biometrika

[Multiple Imputation Strategies for Multiple Group Structural Equation Models](#)

Related research [People also read](#)[Recommended articles](#)[Cited by  
15](#)

## Information for

[Authors](#)[R&D professionals](#)[Editors](#)[Librarians](#)[Societies](#)

## Opportunities

[Reprints and e-prints](#)[Advertising solutions](#)[Accelerated publication](#)[Corporate access solutions](#)

## Open access

[Overview](#)[Open journals](#)[Open Select](#)[Dove Medical Press](#)[F1000Research](#)

## Help and information

[Help and contact](#)[Newsroom](#)[All journals](#)[Books](#)

## Keep up to date

Register to receive personalised research and resources  
by email

[Sign me up](#)