



Structural Equation Modeling: A Multidisciplinary Journal >

Volume 20, 2013 - [Issue 3](#)

544 | 15 | 0
Views | CrossRef citations to date | Altmetric

Original Articles

Investigating Factorial Invariance of Latent Variables Across Populations When Manifest Variables Are Missing Completely

Keith F. Widaman, Kevin J. Grimm, Dawnté R. Early, Richard W. Robins & Rand D. Conger

Pages 384-408 | Published online: 22 Jul 2013

🗨️ Cite this article 🔗 <https://doi.org/10.1080/10705511.2013.797819>

Sample our
Mathematics & Statistics
Journals



>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

📄 Full Article

🖼️ Figures & data

📖 References

🗨️ Citations

📊 Metrics

📄 Reprints & Permissions

Read this article

🔗 Share

Abstract

Difficulties arise in multiple-group evaluations of factorial invariance if particular manifest variables are missing completely in certain groups. Ad hoc analytic alternatives can be used in such situations (e.g., deleting manifest variables), but some common approaches, such as multiple imputation, are not viable. At least 3 solutions to this problem are viable: analyzing differing sets of variables across groups, using pattern mixture approaches, and a new method using random number generation. The latter solution, proposed in this article, is to generate pseudo-random normal deviates for all observations for manifest variables that are missing completely in a given sample and then to specify multiple-group models in a way that respects the random

nature of these values. An empirical example is presented in detail comparing the 3 approaches. The proposed solution can enable quantitative comparisons at the latent variable level between groups using programs that require the same number of manifest variables in each group.

Keywords:

- confirmatory factor analysis
- factorial invariance
- missing data
- structural equation modeling

Notes

- ¹EQS ([Bentler, 2006](#)) scripts and output files for all models presented in this article are available on request from the first author.
- ²Mplus ([Muthén & Muthén, 2007](#)) scripts and output files for all pattern mixture models presented in this article are available on request from the first author.
- ³Analysis scripts and output files for all models presented in this article using the random data approach are available on request from the first author.

Related Research Data

- [Maximum Likelihood from Incomplete Data Via the EM Algorithm](#)
Source: Journal of the Royal Statistical Society Series B (Statistical Methodology)
- [Integrative data analysis through coordination of measurement and analysis protocol across independent longitudinal studies.](#)
Source: Psychological Methods
- [On structural equation modeling with data that are not missing completely at random](#)
Source: Psychometrika
- [Exploring the measurement invariance of psychological instruments: Applications in the substance use domain.](#)
Source: Unknown Repository
- [Inference and Missing Data](#)
Source: Biometrika
- [Multiple Imputation Strategies for Multiple Group Structural Equation Models](#)

Related research

People also read

Recommended articles

Cited by
15

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by email



Sign me up

