



Journal of Liquid Chromatography & Related Technologies >

Volume 30, 2007 - [Issue 12](#)

69 | 18 | 0
Views | CrossRef citations to date | Altmetric

Original Articles

Determination of the Lipophilicity Parameters R_{M0} and $\log P$ of New Azaphenothiazines by Reversed-Phase Thin-Layer Chromatography[†]

Beata Morak, Małgorzata Nowak, & Krystian Pluta

Pages 1845-1854 | Received 10 Dec 2005, Accepted 20 Nov 2006, Published online: 07 May 2007

Cite this article <https://doi.org/10.1080/10826070701360749>

Sample our
Physical Sciences
Journals



>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

Read this article

Share

Abstract

The lipophilicity parameters (R_{M0} and $\log P_{TLC}$) of three types of azaphenothiazines 1–3 were determined by reversed-phase thin-layer chromatography on RP-18 silica plates with acetone-aqueous TRIS (tris(hydroxymethyl)aminomethane) buffer as the mobile phase. The R_M values were linearly dependent on the concentration of acetone, and extrapolated to 0% of acetone, gave the lipophilicity parameter R_{M0} . The parameter R_{M0} and specific hydrophobic surface area b were significantly intercorrelated showing a congeneric class of azaphenothiazines 1–3. The parameter $\log P_{TLC}$ was determined from the R_{M0} values by use of a calibration curve obtained for five standards. The determined parameters were discussed in the terms of structure lipophilicity relationships and compared with data obtained from seven calculation programs.

Keywords:

Lipophilicity parameters

R_{M0}

LogP

Azaphenothiazines

Reversed-phase

TLC

Notes

†Part XCI in the series of Azinyl Sulfides.

Related Research Data

[Calculation Procedures for Molecular Lipophilicity: a Comparative Study](#)

Source: Quantitative Structure-Activity Relationships

[Calculating log Poct from structures](#)

Source: Chemical Reviews

[SYNTHESIS AND PROPERTIES OF 14-SUBSTITUTED 1,4- THIAZINODIQUINOLINES](#)

Source: Phosphorus Sulfur and Silicon and the Related Elements

[Quantitative correlations between albumin binding constants and chromatographic \$R_M\$ values of phenothiazine derivatives](#)

Source: Journal of Medicinal Chemistry

[Lipophilicity of Some N- and O-Substituted Alkanoic Acids of 1,2-Benzisothiazol-3\(2H\)-one Determined by Reversed-Phase Thin Layer Chromatography](#)

Source: Journal of Liquid Chromatography & Related Technologies

[Synthesis of new pentacyclic diquinothiazines](#)

Source: Journal of Heterocyclic Chemistry

[Relationship between the ability of some neuroleptics to enhance striatal](#)



Related research

Recommended articles

Cited by
18

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright © 2026 Informa UK Limited [Privacy policy](#)

[Cookies](#) [Terms & conditions](#) [Accessibility](#)

 Taylor and Francis
Group

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG