



66 | 17 | 0
Views | CrossRef citations to date | Altmetric

Original Articles

Determination of the Lipophilicity Parameters R_{M0} and $\text{Log}P$ of New Azaphenothiazines by Reversed-Phase Thin-Layer Chromatography[†]

Beata Morak, Małgorzata Nowak, & Krystian Pluta

Pages 1845-1854 | Received 10 Dec 2005, Accepted 20 Nov 2006, Published online: 07 May 2007

Cite this article <https://doi.org/10.1080/10826070701360749>

Sample our
Physical Sciences
Journals



>> **Sign in here** to start your access
to the latest two volumes for 14 days

We Care About Your Privacy

We and our 911 partners store and access personal data, like browsing data or unique identifiers, on your device. Selecting I Accept enables tracking technologies to support the purposes shown under we and our partners process data to provide. Selecting Reject All or withdrawing your consent will disable them. If trackers are disabled, some content and ads you see may not be as relevant to you. You can resurface this menu to change your choices or withdraw consent at any time by clicking the Show Purposes link on the bottom of the webpage. Your choices will have effect within our Website. For more details, refer to our Privacy Policy. [Here](#)

We and our partners process data to provide:

Use precise geolocation data. Actively scan device

I Accept

Reject All

Show Purpose

Keywords:

Lipophilicity parameters

 R_{M0}

LogP

Azaphenothiazines

Reversed-phase

TLC

Notes

[†]Part XCI in the series of Azinyl Sulfides.

Related Research Data

Calculation Procedures for Molecular Lipophilicity: a Comparative Study

Source: Quantitative Structure-Activity Relationships

Calculating log Poct from structures

Source: Chemical Reviews

SYNTHESIS AND PROPERTIES OF 14-SUBSTITUTED 1,4- THIAZINODIQUINOLINES

Source: Phosphorus Sulfur and Silicon and the Related Elements

Quantitative correlations between albumin binding constants and chromatographic R_M values of phenothiazine derivatives

Source

Lipone [3(2H)-

Source

Synth

Source

Relat

[3]



Synth

selen

Source

Deter

phase

ones

Source



Multivariate analysis of experimental and computational descriptors of molecular lipophilicity.

Source: Journal of Computer-Aided Molecular Design

UNEXPECTED SIMPLE ROUTE TO NOVEL DIPYRIDO-1,4-THIAZINE SYSTEM

Source: Heterocyclic Communications

AzinyI sulfides. L. 14-Methyl-1,4-thiazino[2,3-c;6,5-c']diquinoline

Source: Acta Crystallographica Section C Crystal Structure Communications

Octanol: Water partition coefficients (P): Measurement, estimation, and interpretation, particularly for chemicals with $P > 10^5$

Source: Ecotoxicology and Environmental Safety

AzinyI sulfides. L.


Source: Acta Crystallographica Section C Crystal Structure Communications

A new method for the estimation of partition coefficient

Source: Journal of the American Chemical Society

Determination of lipophilicity by means of reversed-phase thin-layer chromatography

Source: Journal of Chromatography A

Linking provided by 

Optimize your research
with CHEMPLANE

×





AI ASSISTANT
for Chemistry

TRY IT FREE
FOR 7 DAYS

**OPTIMIZE
YOUR
RESEARCH
WITH
CHEMY
LANE**

Related research 



Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email

 Sign me up



Copyright © 2024

Accessibility

Registered in England
5 Howick Place

Wiley and John Wiley & Francis Group
are trademarks of John Wiley & Sons, Inc., a
Wiley brand business

