



75 | 16 | 3
Views | CrossRef citations to date | Altmetric

Original Articles

A Semi-Explicit Approach to Canary Swaptions in HJM One-Factor Model

Marc Henrard

Pages 1-18 | Received 16 Apr 2004, Published online: 02 Feb 2007

Cite this article <https://doi.org/10.1080/13504860500117602>

Sample our
Economics, Finance,
Business & Industry Journals
>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

Full Article

Figures & data

References

Citations

Metrics

Reprints & Permissions

Read this article

Share

Abstract

Leveraging the explicit formula for European swaptions and coupon-bond options in the HJM one-factor model, a semi-explicit formula for 2-Bermudan options (also called Canary options) is developed. The European swaption formula is extended to future times. So equipped, one is able to reduce the valuation of a 2-Bermudan swaption to a single numerical integration at the first expiry date. In that integration the most complex part of the embedded European swaptions valuation has been simplified to perform it only once and not for every point. In a special but very common in practice case, a semi-explicit formula is provided. Those results lead to a significantly faster and more precise implementation of swaption valuation. The improvements extend even more favourably to sensitivity calculations.

Keywords:

Bermudan swaption

HJM one-factor model

Hull-White model

explicit formula

numerical integration

Acknowledgments

The views expressed here are those of the author and not necessarily those of the Bank for International Settlements. The author wishes to thank the referee and his colleagues for valuable comments on previous versions of the paper. He also wishes to thank an anonymous and careful reader for numerous drafting suggestions.

Notes

1. Bounded is too strong for the proof we use, some L^1 and L^2 conditions are enough, but as all the examples we present are bounded, we use this condition for simplicity.
 2. See Hunt and Kennedy ([2000](#)) for the definition of a numeraire pair. Note that here we require that the bonds of all maturities are martingales for the numeraire pair (N, N) .
 3. Matlab code available from the author.
 4. There is nothing special about that date, except it is my sister's birthday!
 5. As the second step is shorter (6m), the distance between points is also smaller and more than $4n+1$ final points are used.
 6. It took around four hours on my computer to run the (non-optimized) code to compute 3×401 yield curves and the prices for the 4 implementations using 200 steps precisions. As can be inferred from figure [2](#), most of the time was devoted to the tree computations.
-

Related research

People also read

Recommended articles

Cited by
16

Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2025 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)

 Taylor and Francis Group

Registered in England & Wales No. 01072954
5 Howick Place | London | SW1P 1WG