



Q

Home ► All Journals ► Mathematics, Statistics & Data Science ► Applied Mathematical Finance ► List of Issues ► Volume 12, Issue 4 ► Calibration of the SABR Model in Illiqui

Applied Mathematical Finance > Volume 12, 2005 - Issue 4

618 37 0 Views CrossRef citations to date Altmetric

Original Articles

Calibration of the SABR Model in Illiquid Markets Graeme West

Pages 371-385 | Received 14 Jul 2004, Published online: 17 Feb 2007

G Cite this article **I** https://doi.org/10.1080/13504860500148672



Abstract

Recently the SABR model has been developed to manage the option smile which is observed in derivatives markets. Typically, calibration of such models is straightforward as there is adequate data available for robust extraction of the parameters required asinputs to the model. The paper considers calibration of the model in situations where input data is very sparse. Although this will require some creative decision making, the algorithms developed here are remarkably robust and can be used confidently for mark to market and hedging of option portfolios.

Keywords:

Acknowledgment

Thanks to Adam Myers of ABSA Bank, Leon Sanderson of Nedcor Bank, Arthur Phillips of RiskWorX, and Patrick Hagan of Bloomberg's.

Notes

1. Some care needs to be taken with machine precision issues here. One can have that $z\approx 0$ and $\chi(z) = 0$ to double precision. This needs to be trapped, and the limit result invoked, again putting .

2. Although significantly, not all. Real Africa Durolink, a smaller bank, but major player in the equity derivatives market, failed within days of the introduction of the skew, as they were completely unprepared for the dramatic impact the new methodology would have on their margin requirements. See West (2005, Section 13.4).

3. The only difference here is that we do not make the assumption of zero means, which we do when using returns to calculate volatilities. The implementation is elementary.

4. When there are three real roots, they are of the order of -1000, 1 and +1000. So we take the root of order 1.

5. Of course, some experimentation with the choice of the weight determined by the quantum is necessary. One could choose Q 2 for example, or indeed any positive weight. There will be no requirement for any smoothness of the weight in what follows.

6. Note that the matrix will almost certainly not be of size greater than 5×5 .

7. Of course, to very high precision they are always changing. Here we mean that they are unchanged up to the (fairly high) precision that we chose in the Nelder–Mead algorithm.

Related research 🕯

People also read

Recommended articles

Cited by 37

Information for	Open access
Authors	Overview
R&D professionals	Open journals
Editors	Open Select
Librarians	Dove Medical Press
Societies	F1000Research
Opportunities	Help and information
Reprints and e-prints	Help and contact
Advertising solutions	Newsroom
Accelerated publication	All journals
Corporate access solutions	Books

Keep up to date

Register to receive personalised research and resources by email





Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions

Taylor & Francis Group an informa business



Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG