







Q

Home ► All Journals ► Mathematics, Statistics & Data Science ► Applied Mathematical Finance ► List of Issues ► Volume 12, Issue 4 ► Calibration of the SABR Model in Illiqui ....

### Applied Mathematical Finance >

Volume 12, 2005 - Issue 4

638 37 0

Views CrossRef citations to date Altmetric

**Original Articles** 

# Calibration of the SABR Model in Illiquid Markets

Graeme West

Pages 371-385 | Received 14 Jul 2004, Published online: 17 Feb 2007

Sample our
Economics, Finance,
Business & Industry Journals
>> Sign in here to start your access
to the latest two volumes for 14 days









Metrics



Read this article

Share

## Abstract

Recently the SABR model has been developed to manage the option smile which is observed in derivatives markets. Typically, calibration of such models is straightforward as there is adequate data available for robust extraction of the parameters required asinputs to the model. The paper considers calibration of the model in situations where input data is very sparse. Although this will require some creative decision making, the algorithms developed here are remarkably robust and can be used confidently for mark to market and hedging of option portfolios.

Keywords:

SABR model

equity derivatives

volatility skew calibration

illiquid markets

# Acknowledgment

Thanks to Adam Myers of ABSA Bank, Leon Sanderson of Nedcor Bank, Arthur Phillips of RiskWorX, and Patrick Hagan of Bloomberg's.

## Notes

- 1. Some care needs to be taken with machine precision issues here. One can have that  $z\approx 0$  and  $\chi(z)=0$  to double precision. This needs to be trapped, and the limit result invoked, again putting.
- 2. Although significantly, not all. Real Africa Durolink, a smaller bank, but major player in the equity derivatives market, failed within days of the introduction of the skew, as they were completely unprepared for the dramatic impact the new methodology would have on their margin requirements. See West (2005, Section 13.4).
- 3. The only difference here is that we do not make the assumption of zero means, which we do when using returns to calculate volatilities. The implementation is elementary.
- 4. When there are three real roots, they are of the order of -1000, 1 and +1000. So we take the root of order 1.
- 5. Of course, some experimentation with the choice of the weight determined by the quantum is necessary. One could choose Q<sup>2</sup> for example, or indeed any positive weight. There will be no requirement for any smoothness of the weight in what follows.
- 6. Note that the matrix will almost certainly not be of size greater than  $5 \times 5$ .
- 7. Of course, to very high precision they are always changing. Here we mean that they are unchanged up to the (fairly high) precision that we chose in the Nelder-Mead algorithm.

Information for

**Authors** 

**R&D** professionals

**Editors** 

Librarians

**Societies** 

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

**Open Select** 

**Dove Medical Press** 

F1000Research

Help and information

Help and contact

Newsroom

All journals

**Books** 

#### Keep up to date

Register to receive personalised research and resources by email



Sign me up











Accessibility



Copyright © 2025 Informa UK Limited Privacy policy Cookies Terms & conditions



Registered in England & Wales No. 01072954 5 Howick Place | London | SW1P 1WG